

Original article:

PROTECTIVE EFFECT OF AQUEOUS EXTRACT OF THE RHIZOMES OF *SANSEVIERIA LIBERICA* GÉRÔME AND LABROY ON CARBON TETRACHLORIDE INDUCED HEPATOTOXICITY IN RATS

Jude Chigozie Ikewuchi^{1,*}, Catherine Chidinma Ikewuchi¹, Ngozi Mercy Igboh², Telema Mark-Balm¹

¹ Department of Biochemistry, Faculty of Science, University of Port Harcourt, P.M.B. 5323, Port Harcourt, Nigeria

² Department of Biochemistry, Faculty of Science, Abia State University, P.M.B. 2000, Uturu, Nigeria

* Corresponding author: E-mail: ecoli240733@yahoo.com; Tel.: +2348033715662

ABSTRACT

The ability of an aqueous extract of the rhizomes of *Sansevieria liberica* to protect against carbon tetrachloride induced liver injury was investigated in Wistar albino rats. The carbon tetrachloride was prepared 1:5 (v:v) in olive oil, and administered subcutaneously at 1 mL/kg body weight. The extract was administered to both normal and carbon tetrachloride treated rats at 100, 200 and 300 mg/kg. On gas chromatographic analysis of the extract, twenty nine known flavonoids were detected, consisting mainly of 31.94 % apigenin, 20.66 % quercetin, 11.28 % kaempferol, 5.99 % naringenin, 5.83 % (-)-epicatechin, 3.69 % biochanin, 3.58 % (+)-catechin, 2.72 % diadzein, 2.20 % ellagic acid, 2.04 % butein. Compared to test control, the treatment dose dependently produced significantly ($P < 0.05$) lower alkaline phosphatase, aspartate transaminase and alanine transaminase activities. The plasma total bilirubin and total protein levels of the test animals were lower though not significantly. The hepatic histopathological studies showed that carbon tetrachloride caused fatty degeneration of hepatocytes, which was inhibited by pre-treatment with the extract; thus, confirming the results of the biochemical studies. The results of this study indicated that treatment with the plant extracts protects the liver against carbon tetrachloride induced hepatotoxicity. This supports the use of *Sansevieria liberica* in traditional health care for managing liver problems.

Keywords: apigenin, carbon tetrachloride, flavonoids, hepatospecific markers, histopathology, *Sansevieria liberica* Gérôme and Labroy

INTRODUCTION

Liver dysfunction as a result of infection, toxic chemicals, certain drugs and environmental pollutants has been largely increased in the last few decades (Atta et al., 2006). Herbs play a major role in the management of various liver disorders. *Sansevieria liberica* is one of such plants used in African traditional health care for the treatment of liver disorders.

Sansevieria liberica (Family Agavaceae or Ruscaceae or Dracaenaceae), is one of the bowstring hemp species (Evans, 2005), with concave, short petioled leaves that are in part transversely banded with light and dark green, or may be linearly striated with whitish to light green and dark green striations (Reed, 1978). The leaves are very rich in fibers (Ikewuchi et al., 2010a; Osabohien and Egboh, 2008), protein (Ikewuchi et al., 2010a), potassium, calcium, magnesium,

vitamin C, biotin, and riboflavin (Ikewuchi and Ikewuchi, 2009). This plant has long rhizomes with long fibrous roots and a rapid rate of growth, and produces red or orange berry fruits. It is a rather stout herb with several stiff red-margined leaves about 2 feet high arising from the creeping plant, 50-80 cm long inflorescence longer than leaves with abundant white flowers. They are grown as ornamental plants (United States Department of Agriculture, 2008), and are widely distributed throughout the tropics. In Nigeria, the leaves and roots of *Sansevieria liberica* are used in traditional medicine for the treatment of abdominal pain, asthma, colic, diarrhea, eczema, gonorrhoea, hemorrhoid, hypertension, monorrhagia, piles, sexual weakness, wounds, and alleviating the effect snake bite (Adeyemi et al., 2009; Amida et al., 2007; Gill, 1992; Osabohien and Egboh, 2008; Osabohien, 2009). The roots are used for the treatment of convulsion, epilepsy, paralysis, malnutrition, pulmonary troubles, vermifuges, cough and debility. The anti-anaemic (Ikewuchi et al., 2010b) and sedative and anti-convulsant (Adeyemi et al., 2007) activities of the leaves and roots have been reported. In this study, the ability of an aqueous extract of the rhizomes of *Sansevieria liberica*, to protect against carbon tetrachloride induced liver damage was investigated in Wistar albino rats.

MATERIALS AND METHODS

Preparation of plant extract

Samples of fresh *Sansevieria liberica* plants (Figure 1) were procured from horticulturists by Air Force Gate, Aba Road, and at the University of Port Harcourt's Abuja campus, and from behind the Ofrima complex of University of Port Harcourt, all in Port Harcourt, Nigeria. After due identification at the University of Port Harcourt Herbarium, Port Harcourt, the identity was confirmed/authenticated by Dr. Michael C. Dike of Taxonomy Unit, Department of Forestry and Environmental Management, Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria; and Mr.

John Ibe, the Herbarium Manager of the Forestry Department, National Root Crops Research Institute (NRCRI), Umuahia, Nigeria. The rhizomes were removed, rid of dirt, oven dried at 55 °C and ground into powder. The resultant powder was soaked in hot distilled water for 12 h, after which the resultant mixture was filtered and the filtrate (herein referred to as the aqueous extract) was stored in the refrigerator for subsequent use. A known volume of this extract was evaporated to dryness, and the weight of the residue was used to determine the concentration of the filtrate, which was in turn used to determine the dose of administration of the extract. The percentage recovery of the crude extract was 29 %. The resultant residue from the aqueous extract was used for the phytochemical study, in order to determine its flavonoid composition.



Figure 1: *Sansevieria liberica* Gérôme and Labroy

Determination of the flavonoid content of the leaf extract

Calibration, identification and quantification

Standard solutions were prepared in methanol. The linearity of the dependence of response on concentration was verified by regression analysis. Identification was based on comparison of retention times and spectral data with standards. Quantification was performed by establishing calibration curves for each compound determined, using the standards. The chromatogram of the extract is shown in Figure 2.




Figure 2: Chromatogram of the flavonoid composition of an aqueous extract of the rhizomes of *Sansevieria liberica*

Determination of flavonoid composition

The extraction was carried out according to the method of Millogo-Kone et al. (2009). The residue from the aqueous extract above was extracted with methanol and the resultant extract was subjected to gas chromatographic analysis. Chromatographic analyses were carried out on an HP 6890 (Hewlett Packard, Wilmington, DE, USA), GC apparatus, fitted with a flame ionization detector (FID), and powered with HP Chemstation Rev. A 09.01 [1206] software, to quantify and identify compounds. The column was a capillary HP INNOWax Column (30 m × 0.25 mm × 0.25 μm film thickness). The inlet and detection temperatures were 250 and 320 °C. Split injection was adopted with a split ratio of 20:1. Nitrogen was used as the carrier gas. The hydrogen and compressed air pressures were 22 psi and 35 psi. The oven was programmed as follows: initial temperature at 50 °C, first ramping at 8 °C/min for 20 min, maintained for 4 min, followed by a second ramping at 12 °C/min for 4 min, maintained for 4 min.

Experimental design for the hepatoprotective study

Male Wistar albino rats (190-210 g) were collected from the animal house of the Department of Physiology, University of Nigeria, Enugu Campus. Studies were con-

ducted in compliance with applicable laws and regulations for handling experimental animals. The rats were weighed and sorted into eight groups (Table 1) of five animals each, so that their average weights were approximately equal.

Table 1: Experimental design for the hepatoprotective screening

S/N	ID	Treatment
1	Normal control	Olive oil (1 mL/kg) and Normal saline and water
2	Test control	Carbon tetrachloride (1 mL/kg) and water
3	Treatment control I (LRC1)	Olive oil (1 mL/kg) and extract (100 mg/kg)
4	Treatment control II (LRC2)	Olive oil (1 mL/kg) and extract (200 mg/kg)
5	Treatment control III (LRC3)	Olive oil (1 mL/kg) and extract (300 mg/kg)
6	Treatment I (LR1)	Carbon tetrachloride (1 mL/kg) and extract (100 mg/kg)
7	Treatment II (LR2)	Carbon tetrachloride (1 mL/kg) and extract (200 mg/kg)
8	Treatment III (LR3)	Carbon tetrachloride (1 mL/kg) and extract (300 mg/kg)

The animals were housed in plastic cages in the animal house of the Department of Biochemistry, University of Port Harcourt. After a one-week acclimatization period on guinea growers mash (Port Harcourt Flour Mills, Port Harcourt, Nigeria),

the treatment commenced. The extracts were administered orally on daily basis for eight days. The dosages of administration of the extract was adopted and modified from Adeyemi et al. (2007) and Ikewuchi *et al.* (2010b). The carbon tetrachloride was prepared 1:5 (v:v) in olive oil, and administered subcutaneously at 1 mL/kg body weight of carbon tetrachloride, on days 4 and 8. The dosage and method of administration of carbon tetrachloride was adapted from Obi and Uneh (2003), with modification. Twenty four hours after the last administration of carbon tetrachloride, the rats were weighed and anaesthetized by exposure to chloroform. While under anesthesia, they were painlessly sacrificed and blood was collected from each rat into heparin sample bottles, after which the livers were collected and preserved in 10 % formalin, for histological studies. The heparin anticoagulated blood samples were centrifuged at 1000 g for 10 min, after which their plasma was collected and stored for subsequent analysis.

Determination of plasma hepatospecific markers

The plasma activities of alanine transaminase, aspartate transaminase, and alkaline phosphatase, were determined using Randox test kits (Randox Laboratories Ltd., Crumlin, England, UK). The activities of alanine and aspartate transaminases were respectively measured by monitoring at 546 nm the concentrations of pyruvate and oxaloacetate hydrazones formed with 2,4-dinitrophenylhydrazine. The activity of alkaline phosphatase was determined by monitoring the degradation of p-nitrophenylphosphate to p-nitrophenol, at 405 nm.

Plasma total bilirubin and protein concentrations were determined using Randox test kits (Randox Laboratories Ltd., Crumlin, England, UK). The wavelength for the determination of total bilirubin was 578 nm, while that of total protein was 560 nm.

Determination of percentage protection (% protection)

The percentage protection provided by the extract against carbon tetrachloride induced liver damage was calculated using the formula below (Eqn 1) adapted from Al-Qarawi et al. (2004).

Histopathological study

The histopathology study was carried out by Professor S.O. Nwosu, of the Department of Anatomical Pathology, University of Port Harcourt Teaching Hospital. Small pieces of liver tissues were collected in 10 % formalin for proper fixation. These tissues were processed and embedded in paraffin wax. Sections of 5-6 μm in thickness were cut, mounted on slide and stained with hematoxylin and eosin. The sections were then examined via light microscopy (Opticphot-2; Nikon, Tokyo, Japan) at 100 \times magnification.

Statistical analysis of data

All values are reported as the mean \pm s.e.m. (standard error in the mean). The values of the various parameters were analyzed for statistical significant differences between the groups, using the Student's *t*-test, with the help of SPSS Statistics 17.0 package (SPSS Inc., Chicago Ill). $P < 0.05$ was assumed to be significant. Graphs were drawn using Microsoft Office Excel, 2010 software.

$$\% \text{ Protection} = \frac{(\text{Parameter}_{\text{Test control}} - \text{Parameter}_{\text{Treatment}})}{\text{Parameter}_{\text{Test control}} - \text{Parameter}_{\text{Control}}} \times 100 \quad [\text{Eqn 1}]$$

RESULTS AND DISCUSSION

Table 2 shows the flavonoid composition of an aqueous extract of the rhizomes of *Sansevieria liberica*. Twenty nine known flavonoids were detected, consisting mainly of 31.94 % apigenin, 20.66 % quercetin, 11.28 % kaempferol, 5.99 % naringenin, 5.83 % (-)-epicatechin, 3.69 % biochanin, 3.58 % (+)-catechin, 2.72 % diadzein, 2.20 % ellagic acid, 2.04 % butein, 1.85 % robinetin, 1.39 % baicalein, 1.32 % (-)-epigallocatechin, 1.10 % genistein and 1.04 % (+)-gallocatechin. Most of these compounds are bioactive, with antineoplastic and anticarcinogenic properties (Dillard and German, 2000; Evans, 2005).

Figures 3-4 show the effects of an aqueous extract of the rhizomes of *Sansevieria liberica* on plasma markers of liver function. The carbon tetrachloride treatment produced significantly higher ($P<0.05$) plasma alkaline phosphatase, alanine and aspartate transaminases activities, and total bilirubin and total protein levels. Plasma alkaline phosphatase, alanine and aspartate transaminases activities, and total bilirubin level of treated animals were significantly lower ($P<0.05$) than test control. The plasma total protein concentrations of the

treated animals (except LR2) were significantly lower ($P<0.05$) than test control.

Table 2: Composition of flavonoid fraction of an aqueous extract of the rhizomes of *Sansevieria liberica*

Compounds	Retention time (min)	Composition (mg/kg)
(+)-Catechin	13.741	548.722
(+)-Gallocatechin	15.033	159.606
Genistein	15.619	168.342
Diadzein	16.040	416.645
Apigenin	16.247	4895.782
Butein	16.673	312.909
Naringenin	16.785	918.626
Biochanin	17.094	565.454
Luteolin	17.363	117.051
Kaempferol	18.495	1729.383
(-)-Epicatechin	19.522	893.058
(-)-Epigallocatechin	20.471	201.668
Quercetin	21.439	3166.743
(-)-Epicatechin-3-gallate	22.604	15.147
(-)-Epigallocatechin-3-gallate	23.117	93.878
Isorhamnetin	34.098	36.090
Robinetin	24.187	282.869
Ellagic acid	24.611	337.707
Myricetin	24.791	103.970
Baicalein	25.695	212.795
Nobiletin	26.293	96.082
Baicalin	26.950	42.729
Silymarin	27.801	12.947
Total		15328.137




Figure 3: Effects of an aqueous extract of the rhizomes of *Sansevieria liberica* on the activities of plasma hepatospecific marker enzymes in normal and carbon tetrachloride treated rats. Values are mean \pm S.D., $n=5$, per group. ^{a,b,c}Values in the same group with different superscripts are significantly different at $P<0.05$.




Figure 4: Effects of an aqueous extract of the rhizomes of *Sansevieria liberica* on the concentrations of plasma hepatospecific marker molecules in normal and carbon tetrachloride treated rats. Values are mean \pm S.D., n=5, per group. ^{a,b,c}Values in the same column with different superscripts are significantly different at $P < 0.05$.

The hepatoprotective activity of an aqueous extract of the rhizomes of *Sansevieria liberica* on carbon tetrachloride-induced hepatotoxicity in Wistar rats is shown in Figure 5. The protection seemed to be dose dependent. The frequency distribution of the effects of an aqueous extract of the rhizomes of *Sansevieria liberica* on the liver histology of normal and carbon tetrachloride treated rats is presented in Figure 6. Sections of the liver samples are shown in Figure 7. Histopathological stud-

ies of liver sections, showed that carbon tetrachloride caused fatty degeneration, necrosis and ballooning degeneration of hepatocytes; while pre-treatment with aqueous extract of the rhizomes of *Sansevieria liberica* exhibited protection, which confirmed the results of the biochemical studies. The results of our study indicated that treatment with the plant extract protects the liver against carbon tetrachloride-induced hepatotoxicity.




Figure 5: Hepatoprotective activity of an aqueous extract of the rhizomes of *Sansevieria liberica* on carbon tetrachloride-induced hepatotoxicity in Wistar rats




Figure 6: Frequency distribution of the effects of an aqueous extract of the rhizomes of *Sansevieria liberica* on the histology of normal and carbon tetrachloride treated rats




Figure 7: Sections (x20) of the liver samples of showing effect of an aqueous extract of the rhizomes of *Sansevieria liberica* on the liver histology of normal and carbon tetrachloride treated rats. A: Section of the liver of rats administered olive oil (1 mL/kg) and treated with water, showing normal cells. B: Section of the liver tissue of rats administered carbon tetrachloride (1 mL/kg) and treated with water, showing fatty change. C: Section of the liver of rats administered olive oil (1 mL/kg) and treated with 100 mg/kg extract, showing normal cells. D: Section of the liver of rats administered olive oil (1 mL/kg) and treated with 200 mg/kg extract, showing normal cells. E: Section of the liver of rats administered olive oil (1 mL/kg) and treated with 300 mg/kg extract, showing normal cells. F: Section of the liver of rats administered carbon tetrachloride (1 mL/kg) and treated with 100 mg/kg extract, showing normal cells. G: Section of the liver of rats administered carbon tetrachloride (1 mL/kg) and treated with 200 mg/kg extract, showing normal cells. H: Section of the liver of rats administered carbon tetrachloride (1 mL/kg) and treated with 300 mg/kg extract, showing focal areas of fatty change.

From Figures 3 and 4 it is evident that the extract was able to reduce all the elevated biochemical parameters due to the hepatotoxin intoxication. The extract showed very significant hepatoprotection against carbon tetrachloride-induced hepatotoxicity in the rats, by reducing plasma

total bilirubin and protein, plasma alkaline phosphatase, alanine and aspartate transaminases levels. The reduction towards the normal value, of the levels of these plasma indices of liver integrity/function is an indication of the ability of the extract to protect normal structural/functional integrity of the

poisoned liver, and also to protect against subsequent carbon tetrachloride hepatotoxicity, enabling regeneration process. A fact that was confirmed by histopathological studies on liver sections: that revealed that the treated animals had normal hepatic cells. This hepatoprotective activity may have been produced via any of the following mechanisms.

Reduced metabolic activation of carbon tetrachloride by cytochrome P450 depresses the initial formation of trichloromethyl free radical, resulting in the diminished initiation of lipid peroxidation (Middleton et al., 2000), and the consequent toxicity of carbon tetrachloride. So, any hepatoprotective drug should be able to inhibit the aromatase activity of cytochrome P450, and thereby, favor liver regeneration. Therefore, it can be suggested that flavonoids in rhizomes of *Sansevieria liberica* (Table 2), could be responsible for its hepatoprotective ability. Flavonoids have been reported to inhibit lipid peroxidation by exerting a membrane-stabilizing action (Middleton et al., 2000) or inhibiting cytochrome P450 aromatase (Kowalska et al., 1990; Middleton et al., 2000).

Many hepatoprotective flavonoids were detected in the aqueous extract of the rhizomes of *Sansevieria liberica*. They include apigenin, quercetin, kaempferol, naringenin and silymarin. The hepatoprotective activities of apigenin (El Alfy et al., 2010; Zheng et al., 2005), quercetin (Chen, 2010; Lee et al., 2003; Mandal and Das, 2005; Mandal et al., 2007; Pavanato et al., 2003; Peres et al., 2000), kaempferol (Oh et al., 2004; Song et al., 2003; Xiong et al., 2000), naringenin (Lee et al., 2004; Pari and Gnanasoundari, 2006) and silymarin (Suja et al., 2004) have all been documented.

Another component of *Sansevieria liberica* that may also have contributed to its hepatoprotective activity is vitamin C, a compound that has been reported to be abundant in *Sansevieria liberica* (Ikewuchi and Ikewuchi, 2009). In vivo studies have indicated that hepatic microsomal drug metabolism decreases in ascorbic acid defi-

ciency and is augmented when high supplements of the vitamin are given to guinea pigs (Burtis and Ashwood, 2001; Sato and Zannoni, 1976). Rikans *et al.* (1978) had also reported that liver cytochrome P450 is significantly reduced in ascorbic acid-deficiency.

CONCLUSION

This study clearly demonstrates that aqueous extract of the rhizomes of *Sansevieria liberica* is an effective agent in the treatment and prevention of carbon tetrachloride-induced hepatic cytotoxicity. The data suggest that the daily oral consumption of the extract was prophylactic to carbon tetrachloride poisoning. This confirms the use of *Sansevieria liberica* in traditional health care for the treatment of liver problems.

REFERENCES

- Adeyemi OO, Yemitan OK, Adebisi OO. Sedative and anticonvulsant activities of the aqueous root extract *Sansevieria liberica* Gerome and Labroy (Agavaceae). J Ethnopharmacol 2007;113:111-4.
- Adeyemi OO, Akindele AJ, Ogunleye EA. Evaluation of the antidiarrhoeal effect of *Sansevieria liberica* Gerome and Labroy (Agavaceae) root extract. J Ethnopharmacol 2009;123:459-63.
- Al-Qarawi AA, Mousa HM, Ali BE-DH, Abdel-Rahman H, El-Mougy SA. Protective effect of extracts from Dates (*Phoenix dactylifera* L.) on carbon tetrachloride-induced hepatotoxicity in rats. Internat J Appl Res Vet Med 2004;2:176–80.
- Amida MB, Yemitan OK, Adeyemi OO. Toxicological assessment of the aqueous root extract of *Sansevieria liberica* Gerome and Labroy (Agavaceae). J Ethnopharmacol 2007;113:171-5.

- Atta AH, Nasr SM, Mouneir SM. Potential protective effect of some plant extracts against carbon tetrachloride-induced hepatotoxicity. *Afr J Trad Compl Alt Med* 2006;3(3):1-9.
- Burtis CA, Ashwood ER. *Tietz's fundamentals of clinical chemistry*, Vol 2. Philadelphia: WB Saunders, 2001.
- Chen X. Protective effects of quercetin on liver injury induced by ethanol. *Pharmacogn Mag* 2010;6(22):135–41.
- Dillard CJ, German JB. Phytochemicals: nutraceuticals and human health. *J Sci Food Agric* 2000;80:1744–56.
- El Alfy TS, El Sawi SA, Sleem A, Moawad DM. Investigation of flavonoidal content and biological activities of *Chorisia Insignis* Hbk. Leaves. *Aust J Basic Appl Sci* 2010;4:1334-48.
- Evans WC. A taxonomic approach to the study of medicinal plants and animal-derived drugs. In: Evans WC (ed.): *Trease and Evans pharmacognosy*, 15th ed. (pp. 15-40). India: Elsevier, 2005.
- Gill LS. *Ethnomedical uses of plants in Nigeria*. Benin City, Nigeria: Uniben Press, 1992.
- Ikewuchi CC, Ikewuchi JC. Amino acid, mineral, and vitamin composition of *Sansevieria liberica* Gérôme and Labroy. *Pac J Sci Technol* 2009;10:477-82.
- Ikewuchi CC, Ikewuchi CJ, Ayalogu OE, Onyeike NE. Proximate and phytochemical profile of *Sansevieria liberica* Gérôme and Labroy. *J Appl Sci Environ Mgt* 2010a;14: 103–6.
- Ikewuchi CC, Ikewuchi JC, Onyeike EN, Ayalogu EO. Effect of *Sansevieria liberica* Gérôme and Labroy on plasma chemistry and hematological indices of salt-loaded rats. *Res J Sci Technol* 2010b;2(5):110-4.
- Kowalska MT, Brandt ME, Puett D. Inhibition of cytochrome P-450 aromatase activity by plant extracts. *Planta Med* 1990;56: 675–7.
- Lee E-S, Lee H-E, Shin JY, Yoon S, Moon J-O. The flavonoid quercetin inhibits dimethylnitrosamine-induced liver damage in rats. *J Pharmacy Pharmacol* 2003;55:1169–74.
- Lee M-H, Yoon S, Moon J-O. The flavonoid naringenin inhibits dimethylnitrosamine-induced liver damage in rats. *Biol Pharm Bull* 2004;27(1):72-6.
- Mandal AK, Das N. Sugar coated liposomal flavonoid: a unique formulation in combating carbon tetrachloride induced hepatic oxidative damage. *J Drug Target* 2005;13: 305–15.
- Mandal AK, Das S, Basu MK, Chakrabarti RN, Das N. Hepatoprotective activity of liposomal flavonoid against arsenite-induced liver fibrosis. *J Pharmacol Exp Therap* 2007;320:994–1001.
- Middleton E Jr, Kandaswami C, Theoharides TC. The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease and cancer. *Pharmacol Rev* 2000;52:673-751.
- Millogo-Kone H, Lompo M, Kini F, Asimi S, Guissou IP, Nacoulma O. Evaluation of flavonoids and total phenolic contents of stem bark and leaves of *Parkia biglobosa* (Jacq.) Benth. (Mimosaceae)-free radical scavenging and antimicrobial activities. *Res J Med Sci* 2009;3(2):70-4.
- Obi FO, Uneh E. pH dependent prevention of carbon tetrachloride-induced lipoperoxidation in rats by ethanolic extract of *Hibiscus rosasinensis* petal. *Biokemistri* 2003; 13:42-50.

- Oh H, Kim D-H, Chob J-H, Kim Y-C. Hepatoprotective and free radical scavenging activities of phenolic petrosins and flavonoids isolated from *Equisetum arvense*. J Ethnopharmacol 2004;95:421-4.
- Osabohien E. Effects of epoxidation on the thermal conductivity and equilibrium swelling properties of bowstring hemp fibre natural rubber composites. Adv Nat Appl Sci Res 2009;7:169-76.
- Osabohien E, Egboh SHO. Utilization of bowstring hemp fiber as a filler in natural rubber compounds. J Appl Polymer Sci 2008;107:210-4.
- Pari L, Gnanasoundari M. Influence of naringenin on oxytetracycline mediated oxidative damage in rat liver. Basic Clin Pharmacol Toxicol 2006;98:456-61.
- Pavanato A, Tuñón MJ, Sonia SC, Marroni CA, Llesuy S, González-Gallego J et al. Effects of quercetin on liver damage in rats with carbon tetrachloride-induced cirrhosis. Digest Dis Sci 2003;48:824-9.
- Peres W, Tunon, MJ, Collado PS, Herrmann S, Marroni N, Gonzalez-Gallego J. The flavonoid quercetin ameliorates liver damage in rats with biliary obstruction. J Hepatol 2000;33:742-50.
- Reed DW. Sansevieria - Identification and culture. Louisiana Soc Hort Res 1978; 14(2):30-77.
- Rikans LE, Smith CR, Zannoni VG. Ascorbic acid and cytochrome P-450. J Pharmacol Exp Ther 1978;204:702-5.
- Sato PH, Zannoni VG. Ascorbic acid and hepatic drug metabolism. J Pharmacol Exp Ther 1976;198:295-307.
- Song E-K, Kim J-H, Kim J-S, Cho H, Nan J-X, Sohn D-H et al. Hepatoprotective phenolic constituents of *Rhodiola sachalinensis* on tacrine-induced cytotoxicity in Hep G2 cells. Phytotherapy Res 2003;17:563-5.
- Suja SR, Latha PG, Pushpangandan P, Rajasekharan S. Evaluation of hepatoprotective effects of *Helminthostachys zeylanica* (L) Hook against carbon tetrachloride induced liver damage in Wistar rats. J Ethnopharmacol 2004;92:61-6.
- United States Department of Agriculture, Agricultural Research Service, Germplasm Resources Information Network (GRIN). GRIN taxonomy for plants [database on the Internet] [cited 2008 July 23]. Available from: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?333057>
- Xiong Q, Fan W, Tezuka Y, Adnyana IK, Stampoulis P, Hattori M et al. Hepatoprotective effect of *Apocynum venetum* and its active constituents. Planta Med 2000;66(2): 127-33.
- Zheng Q-S, Sun X-L, Xu B, Li G, Song M. Mechanisms of apigenin-7-glucoside as a hepatoprotective agent. Biomed Environ Sci 2005;18:65-70.