

**Original article:**

**CHRYSEN-2-OL DERIVATIVE FROM WEST INDIAN WOOD NETTLE  
*LAPORTEA AESTUANS* (L.) CHEW INHIBITS OXIDATION  
AND MICROBIAL GROWTH *IN VITRO***

Ganiyat K. Oloyede\*, Martha S. Oyelola

Natural products/Medicinal Chemistry Unit, Department of Chemistry, University of Ibadan, Nigeria

\* corresponding author: Ganiyat Kehinde Oloyede,

E-mail: oloyedegk@gmail.com, Telephone: +234 803 562 2238

**ABSTRACT**

Bio-active compounds present in West Indian Wood Nettle *Laportea aestuans* (L.) Chew (Urticaceae), used in ethno medicine as antioxidant and antimicrobial were studied. The aim of this research work was to isolate and characterize the bio-active compounds in the n-hexane fraction of *L. aestuans*, determine the toxicity and subject it to *in-vitro* antimicrobial and free radical scavenging activities. The chemical constituents were isolated by gradient elution column chromatographic technique and Ultra Violet/visible (UV), Infrared (IR) and Nuclear Magnetic Resonance (NMR) spectroscopies were used for structural elucidation. The free radical scavenging activity of the isolate was assessed using three methods; scavenging effect on 2,2-diphenyl-1-picrylhydrazyl radical (DPPH), hydroxyl radical generated from hydrogen peroxide and ferric thiocyanate method. Antimicrobial screening was done by agar well diffusion method while toxicity was determined by Brine shrimp lethality test.

Structures were proposed for the white crystalline solids isolated; (4*E*)-3,6-dimethylhep-4-en-3-ol (AB) and 1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-((*E*)-4,7-dimethyloct-5-enyl) chrysen-2-ol (AC). Percentage yield of AC was 91.2 and was non-toxic with LC<sub>50</sub> (µg/ml) value of 1581233000.0. AC significantly scavenged free radical at 0.0625 mg/ml in the DPPH (64.73 %) and hydrogen peroxide (99.22 %) tests. It also showed 65.23 % inhibition at 1.0 mg/ml in the ferric thiocyanate test. AC also inhibited microbial growth significantly when compared with gentamicin and tioconazole which are antibacterial and antifungal standards respectively. The presence of Chrysen-2-ol derivative in *L. aestuans* which was non-toxic and possessed significant antimicrobial and antioxidant activities supports its ethno medicinal application.

**Keywords:** *Laportea aestuans*, (4*E*)-3,6-dimethylhep-4-en-3-ol, chrysen-2-ol, toxicity, antimicrobial, antioxidant

**INTRODUCTION**

The potential of plants as a veritable source for pharmaceuticals and other therapeutic materials have been emphasized. Plants with medicinal values often referred to as secondary metabolites have been documented in herbal pharmacopeias, directo-

ries and books (Gbile, 1986; Ghani, 1986; Dokosi, 1998). Secondary plant metabolites also referred to as natural products are renowned for their potent pharmacological activities; as antimicrobial, insecticidal, anticancer, antioxidant, pesticidal, amongst others. Some are odoriferous and find ap-

plication in perfumes and cosmetics, for flavoring foods and drinks and for screening incense and household cleaning products. They include alkaloids, flavonoids, steroids, coumarins, anthocyanins, terpenoids and essential oils (Brinker, 1998; Burkill, 1985; Gbile, 1986; Abu-Rabia, 2005; Gilbert and Martin, 2006). In recent years, there has been an extensive growth in natural product chemistry. This is due to advances in isolation, characterization, identification, purification and synthesis of new natural products. Many of today's medicines are obtained directly from natural sources and a renewed interest in the screening of plants for biological activity is due to the developments of *in vivo* and biochemical screening methods (Kokwaro, 1993; Heywood, 1993).

*Laportea aestuans*, the West Indian wood nettle, is an annual herb of the *Urticaceae* or nettle family. It is native to tropical Africa, although it is now widespread as an introduced species throughout both the western hemisphere and eastern hemisphere tropics and subtropics, including the USA. *Laportea aestuans* (Linn) Chew have the following synonyms; *Urtica aestuans*, *Fleurya aestuans* (Gaud) and *Fleurya aestuans* (Linn.) Miq. It is used as abortifacient, antimicrobial, laxative, in eye treatments and pain-killers. It is also used in treating pulmonary and stomach troubles, diarrhea and dysentery (Chew, 1969; Burkill, 1985; Friis, 1993; Nadine, 2001). *Laportea aestuans* (L.) Chew is recorded as an ethno veterinary species that is used for urinary problems of ruminants in Trinidad and Tobago and also to shorten labour and remove the placenta during childbirth (Lans, 2006, 2007). Epidemiological studies indicated that consumption of *L. aestuans* inhibits the damaging activities of free radicals in human body (Morrison and Twumasi, 2010). Free radicals find their way into the human body via metabolic pathways within body tissues and also from external sources such as food, drugs and pollution from the environment (Miller and

Britigan, 1997). It was also suggested that free radicals, such as superoxide radical ( $O_2^-$ ) and hydroxyl radical ( $OH^\cdot$ ) and non-free radical species, such as hydrogen peroxide ( $H_2O_2$ ) and singlet oxygen ( $^1O_2$ ) within the human body facilitate cellular injury, ageing, development of neurodegenerative and cardiovascular diseases (Stadtman, 1992; Ames et al., 1993; Alan and Miller, 1996; Knight, 1998; Pinzino et al., 1999; Cadenas and Davies, 2000; de la Fuente, 2000; Stanner et al., 2004). Consumption of fruits and vegetables has proven to substantially reduce the risk of cardiovascular diseases and neurodegenerative diseases, including Parkinson's and Alzheimer's diseases (Di Matteo and Esposito, 2003; Bjelakovic et al., 2007). Paul (2008) isolated a mixture of methyl esters of aliphatic acids from *Laportea aestuans*. Also, the essential oil from the plant was found to be dominated by methyl salicylate and had significant antioxidant and antimicrobial activities (Oloyede, 2011) while phytochemical, toxicity, antimicrobial and antioxidant screening of extracts obtained from *Laportea aestuans* (Gaud) had also been investigated (Oloyede and Ayanbadejo, 2013). Based on our findings, there was need to isolate and characterize the bioactive compounds in *Laportea aestuans*. The prevalence of microbial infection and oxidation reaction is also a major driving force for this study. This current study is therefore aimed at isolating and characterizing the plant's secondary metabolite using chromatographic and spectroscopic techniques (UV, IR and NMR). Antioxidant and antimicrobial screening of the isolated pure compound was also carried out and the toxicity level was determined using Brine shrimp larvae eggs.

## MATERIALS AND METHODS

### *Chemicals and reagents*

Dimethylsulphoxide (M&B, England), hydrogen peroxide (Merck, Germany) and 2,2-diphenyl-1-picrylhydrazyl (DPPH), silica gel 60-200 mesh size, ascorbic acid, bu-

tylated hydroxyanisole (BHA), rutin and  $\alpha$ -tocopherol were obtained from Sigma Chemical Co (St Louis, MO). Brine shrimp larvae eggs were obtained from Ocean Star International, Inc. Company, USA. Acetone, chloroform, ethyl acetate, n-hexane, methanol, butanol, chloroform, hydrochloric acid, ammonia solution, naphthol, bismuth nitrate, potassium iodide, sodium hydroxide, copper acetate, NaOH, ferrous chloride, sodium chloride, copper sulphate pentahydrate, ferric chloride, conc. tetraoxosulphate (VI) acid, conc. HCl, ammonia solution, sodium potassium tartarate, linoleic acid, ammonium thiocyanate, ethanol, hydrochloric acid, potassium chloride, glacial acetic acid, disodium hydrogen phosphate and dihydrogen potassium phosphate were all BDH general purpose chemicals and solvents were distilled prior to use.

#### **Equipment and apparatus**

Rotavapor R110 (Buchi, England), Soxhlet apparatus, Ohaus analytical balance (USA), Steam Bath (Gallenkamp), silica gel GF<sub>254</sub> (precoated aluminium sheets - Merck Germany), pH meter (Jenway model), Gallenkamp melting point apparatus, UV-Visible spectrophotometer (UVD-2960 model equipped with a UVWIN software version LABOMED INC, USA). IR Spectrophotometer (Perkin Elmer FT-IR system, Spectrum BX model of IR Spectrophotometer equipped with a PC and a Printer, England), Varian-Mercury NMR Spectrophotometer operating at 200 MHz for H and 50 MHz for C nuclei.

#### **Plant collection and identification**

Fresh leaves of *L. aestuans* weighing 13.4 kg were collected from the Botanical Garden, University of Ibadan, Nigeria. Identification of sample was done at the herbarium of Botany Department of the same University in January, 2011, by Mr. Owolabi and at the Forestry Research Institute of Nigeria (FRIN) by Mr. Afilaka.

#### **Test organisms**

*Escherichia coli*, *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Klebsiellae pneumoniae*, *Salmonellae typhi*, *Candida albicans*, *Aspergillus flavus* and *Fusarium moniliformes* (Microorganisms were collected from the stock of the Department of Microbiology, Faculty of Science of University of Ibadan). The test organisms were maintained on nutrient agar slopes and kept in a refrigerator at 4 °C. Aliquots of nutrient broth (100 ml) were inoculated with the culture of test microorganisms using a loop and then incubated at 37 °C for 24 h.

#### **Reference standards**

Gentamicin (80 µg/ml) and nystatin (80 µg/ml) were standards for bacteria and fungi respectively for the antimicrobial screening. Ascorbic acid, butylated hydroxyanisole (BHA),  $\alpha$ -tocopherol and rutin were used for antioxidant activity. Dimethylsulfoxide (DMSO) was used for cytotoxicity studies.

#### **Sample preparation**

Whole plant of *L. aestuans* was collected, sorted to remove dirt, weighed and air dried for 3 weeks until the weight was constant and then pulverized using mill machine at the Mechanical Department of the Faculty Engineering of the University of Ibadan. The pulverized sample (2.355 kg) was kept for further analysis.

#### **Extraction / purification procedure**

Hot extraction was carried out by the use of improvised Soxhlet apparatus. The ground plant sample (2.35 kg) was packed and 5 Lt hexane was poured over the packed sample in the aspirator bottle. The extraction process continued until a colorless or clear hexane was observed indicating the end of the extraction. The extracts were collected and concentrated with the aid of a Buchi rotavapor and stored in a desiccator prior to further analysis. The percentage yield of the extract was determined

by the percent recovery technique. Thin Layer Chromatography (TLC) was employed using silica gel 60 F<sub>254</sub> precoated plates and solvent system: hexane: ethyl acetate (2:5), (2:7), (3:1), (4:1), (5:1), (6:1), (10:1), (15:1). TLC was carried out on the crude extract to estimate the number of components. One drop of glacial acetic acid was added to each solvent system to aid distinct separation of spots into their different components. After the development, the chromatogram was then allowed to dry before it was transferred into the iodine tank. Initial spotting of the crude extract obtained on TLC plates indicated more than one components, hence the need for proper separation with a column chromatograph. Liquid column chromatograph used to separate non-volatile or thermally stable compounds was used to separate the components of the crude extract. 3 cm x 100 cm column size and a silica gel of mesh size 60-200 were used. Silica gel (180 g) was made into slurry by adding the solid adsorbent to a quantity of packing solvent (hexane). The slurry was swirled until it was homogeneous and relatively free of entrapped air. Prepared slurry was slowly poured into the column with the tap opened such that it gently settled uniformly. The column was constantly tapped to get rid of any trapped air bubbles and also ensure a leveled layer of the adsorbent. 6 g of crude extract was pre-adsorbed on silica gel. Hexane was used to wash the packed column prior to analysis. This pre-adsorbed sample was allowed to dry before loading onto the column. The ratio of the silica gel to that of extract was 30:1. The loaded sample was then covered with pure white sand and a small piece of cotton wool. The top of the column was usually filled with enough solvent/mixture of solvents to avoid cracking. The elution started with 100 % non-polar hexane after which the polarity of the solvent(s) was increased gradually with EtOAc until 100 % in a gradient elution chromatographic technique. The solvent system used in eluting the column was as follows: hex-

ane 100 %, ethyl acetate/hexane 5 %, 10 %, 15 %, 20 %, 25 %, 40 %, 50 %, ethyl acetate 100 %. A total of 143 eluents of 100 ml each were collected. TLC of the eluents collected was also carried out using the same solvent system employed in the column chromatography. This was done to know the purity of each eluent, ascertain the number of components in it and also to pool together similar eluents which contain the same component based on their R<sub>f</sub> values. Fractions 38 and 39 and 53 to 60 which were solids obtained from column chromatography were dissolved in hot ethyl acetate and hot acetone respectively and allowed to cool in ice bath in order to form precipitated solutions. The precipitates formed on cooling were separately filtered by vacuum filtration. The crystals were tagged AB and AC. The crystals were left at room temperature for three days in order to dry them. Melting point of pure compounds was determined with a Gallenkamp melting point apparatus. Spectroscopic analysis, UV-Visible, Infra-red and NMR spectrometry was used to characterize the compounds. The UV/Visible Spectra of 0.01 % w/v of the fractions were determined with the aid of UV-Visible spectrophotometer (UVD-2960 model equipped with a UVWIN software version LABOMED Inc., USA). V<sub>max</sub> (cm<sup>-1</sup>) from IR data also confirmed the structures. The Infrared spectrum of the pure compound was determined using a Perkin Elmer FT - IR Spectrophotometer, England). The KBr disc method was used for the preparation of the sample. The relative strength and position of all absorption in the infrared region was determined. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of the pure compounds were determined using a 200 MHz machine for 10 % (w/v) solutions in deuteriochloroform. Pulse irradiation technique employed was FT NMR at ambient temperature. Solid AC with higher yield and promising antioxidant activity in the preliminary screening was investigated for antioxidant and antimicrobial activity and its toxicity was also determined

## Antioxidant activities of AC

### Scavenging effect on DPPH

2, 2-diphenyl-1-picrylhydrazyl radical (DPPH), a stable radical decolorizes in the presence of a free radical scavenger and decreases in absorption in a UV/Visible spectrophotometer when measured 10 min later against that of the control at 517 nm. Antioxidant activity of solid AC and antioxidant standards butylated hydroxyanisole (BHA), rutin,  $\alpha$ -tocopherol and ascorbic acid were determined using the DPPH free-radical scavenging method. A 100  $\mu$ m solution of DPPH was prepared by dissolving 3.94 mg in 100 ml methanol. To 3.0 ml of the methanol solution of DPPH was added 0.5 ml of the samples with doses ranging from 1.0 - 0.0625 mg/ml. Average results from triplicate analysis were recorded and the percentage inhibition of DPPH discoloration was calculated as % inhibition =  $\{(A_{DPPH} - A_S)/A_{DPPH}\} \times 100$ . Where  $A_S$  is the absorbance of the sample solution and  $A_{DPPH}$  is the absorbance of the DPPH solution (Hatano et al, 1988; Gulcin et al, 2002, Pourmorad, 2006; Mutee, 2010).

### Scavenging effect on hydrogen peroxide

Solid AC and standards butylated hydroxyanisole (BHA), ascorbic acid and  $\alpha$ -tocopherol at the following concentrations; 1.0 - 0.0625 mg/ml were each added to a solution of 2 mM hydrogen peroxide, prepared in phosphate buffered-saline (PBS) pH 7.4. Decrease in absorbance of  $H_2O_2$  was determined in a UV/Visible spectrophotometer at 285 nm after 10 min against a blank solution containing the sample in PBS without  $H_2O_2$ . Average result from triplicates analysis was recorded and was used to calculate percentage inhibition (Sies, 1997; Soares et al., 1997, Oloyede and Farombi, 2010).

### Antioxidant activity by ferric thiocyanate method

Ferric thiocyanate method which is another antioxidant screening technique was used to screen compound AC obtained from *L. aestuens* and standards for antioxidant

activity. Various concentrations (1.0 - 0.0625 mg/ml) of sample were prepared from a stock solution containing 10 mg of crystal in 99.5 % of ethanol. A mixture of 4 ml of 0.05 M phosphate buffer (pH 7.0), water (2 ml), sample (2 ml) and 2.0 ml of 2.51 % linoleic acid in 99.5 % ethanol were placed in a vial with a screw cap and placed in the dark in an oven at 6 °C. To 0.1 ml of this sample solution was added 10 ml of 75 % ethanol, 0.1 ml of 30 % ammonium thiocyanate and 0.1 ml of  $2 \times 10^{-2}$  M ferrous chloride in 3.5 % hydrochloric acid, the absorbance of the red color which developed was measured at 500 nm after 3 min. Standards were also subjected to the same experiment. Average result from triplicates analysis was recorded and inhibition of lipid peroxidation in percentage was calculated from this equation:

$$\% \text{ Inhibition} = 1 - (A1/A2) \times 100.$$

Where A1 was the absorbance of the sample and A2 was the absorbance of control reaction (Oloyede et al., 2010).

## Toxicity analysis

### Brine shrimp lethality test

The toxicity of compound AC to living organisms was predicted using the Brine shrimp lethality test (BST). The shrimp's eggs were hatched in sea water for 48 h at room temperature. The *nauplii* (harvested shrimps) were attracted to one side of the vials with a light source. Solutions of the sample were made in DMSO, at varying concentrations (1000, 100 and 10  $\mu$ g/ml) and incubated in triplicate vials with the brine shrimp larvae. Ten brine shrimp larvae were placed in each of the triplicate vials. Control brine shrimp larvae were placed in a mixture of sea water and DMSO only. The vials were examined against a lighted background after 24 h and the average number of larvae that survived in each vial was determined. The concentration at fifty percent mortality of the larvae ( $LC_{50}$ ) was determined using the Finney computer programme (Meyer et al., 1982; Falope et al., 1993; Keddy et al., 1995).

### Antimicrobial screening

#### Agar diffusion: pour plate for bacteria

The following organisms; *Escherichia coli*, *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Klebsiellae pneumoniae* and *Salmonellae typhi* were used. Preparation of culture was done overnight and 0.1 ml of each of the organism was taken into 9.9 ml of Sterile Distilled Water (SDW) to give 10 ml of 1:100 ( $10^2$ ) dilution. A 0.2 ml of 1:100 dilutions was taken into the prepared molten Nutrient Agar (NA) at 45 °C and was aseptically poured into the sterile plates and allowed to set on the bench for 45 min. A sterile cork-borer was used to create wells inside the set plates and different prepared concentrations of the sample as well as the positive and negative controls were introduced using syringes. The positive control for bacteria was gentamicin (80 µg/ml). The plates were left on the laboratory bench for two h before incubation at 37 °C for 24 h. Clear zones of inhibition were observed and the diameters were measured in millimeter (mm) using a transparent well-calibrated ruler. The average of triplicate readings were calculated and recorded (Schmidt, 2006; Duraipandian et al., 2006).

#### Agar diffusion: surface plate for fungi

*Candida albicans*, *Aspergillus flavus* and *Fusarium moniliformes* were the fungal strains used in this experiment. Molten sterile Sabouraud Dextrose Agar (SDA) was poured aseptically into the sterile plates and allowed to cool for 45 min. A 0.2 ml of 1:100 dilution of the organism was spread on the surface using a sterile spreader and a sterile cork-borer was used to create wells inside the plates. The same procedure described for anti-bacterial activity above was followed from this stage. The positive control for the fungi was tioconazole (80 µg/ml). All the plates for the fungi were incubated at 28 °C for 48 h. The clear zones of inhibition were observed and recorded using the same method as described for bacteria (Bayer et al., 1986).


### Data analysis

The mean  $\pm$  SEM was calculated for each parameter and was analyzed separately using ANOVA followed by Dunnett's 't' test. Toxicity results were analyzed using the Finney computer programme.

## RESULTS

Thin Layer Chromatographic analysis (TLC) of the crude extract gave five components with  $R_f$  values 0.39, 0.46, 0.69, 0.84 and 0.94. Column chromatography gave 143 fractions and were bulked based on similarity of their  $R_f$  values as A<sub>1</sub> (fractions 1-20), AA (fractions 21-25), A<sub>2</sub> (Fractions 26-37), AB (fractions 38 and 39), A<sub>3</sub> (fractions 40-52) and AC (fractions 53-60). The remaining fractions 61-76, 77-78, 79-81, 82-89, 90-106, 107-118, 119-125, 126-128, 129-133, 134-140, 141-143 were also pooled accordingly. AA, A<sub>1</sub>, A<sub>2</sub> and A<sub>3</sub> were oily substances while AB and AC were crystalline solids. They all gave single spot in TLC analysis. Fraction AB (yield; 77.7 %) and AC (yield; 91.2 %) were further purified by crystallization. Fraction AB which was light brown solid dissolved in hot ethyl acetate and was allowed to cool in ice bath. The precipitate formed was filtered off giving white solid substance. Isolate AC, a greenish needle - like crystalline solid was dissolved in hot acetone and allowed to cool in ice bath. The precipitate formed was filtered and a white crystalline solid was obtained. Gallenkamp melting point apparatus was used to determine the melting point of the solids. Melting points of AB and AC were 76 °C-78 °C and 131 °C-132 °C respectively. Structures were proposed for solids AB and AC based on the spectra data obtained from UV/visible, infrared (IR), Proton and Carbon 13 Nuclear Magnetic Resonance (<sup>1</sup>H and <sup>13</sup>C NMR) spectroscopies and were in complete agreement with literature values.

Structures of the compounds reported here are as shown in Figure 1.



**Figure 1:** Structures of (4E)-3,6-dimethylhept-4-en-3-ol (AB) and 1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-((E)-4,7-dimethyloct-5-enyl)chrysen-2-ol (AC)

(4E)-3,6-dimethylhept-4-en-3-ol (**AB**): Molecular formulae  $C_9H_{18}O$ , white solid substance. Yield: (8.3 mg) 77.7%. M.pt: 76<sup>o</sup>C-78<sup>o</sup>C. Rf: 0.69 (EtOAc-Hexane, 1:4). UV [EtOH] nm (log  $\epsilon$ ): 207.78 (3.2508) and 254.48 (0.78588). IR (KBr)  $V_{max}$   $cm^{-1}$ : 3432 (O-H), 2908.43 (C-H), 1635.03 (C=C), 1401.95, 1257.74, 1055.39, 771.20, 567.75 (fingerprint region, C-H aliphatic out of plane bending, C-O, C-C etc); <sup>1</sup>H NMR (200 MHz;  $CDCl_3$ ):  $\delta$  3.75 (1H, m, O-H and 2H, C=C-H overlap), 1.6 (2H, m,  $CH_2$ ), 1.4 (10H, m,  $CH_3$ -), 0.9 (3H, t,  $CH_3$ ). <sup>13</sup>C-NMR (50 MHz;  $CDCl_3$ ):  $\delta$  76.374 (C-3, C-O), 63.122 (C-H, alkene C-4), 32.812 (C-5, C-H, alkene), 31.934 (C-6, C-H), 29.708 (C-2,  $CH_2$ ), 29.371 (C-9,  $CH_3$ ), 25.740 (C-7,  $CH_3$ ), 22.694 (C-8,  $CH_3$ ), 14.143 (C-1,  $CH_3$ ).

1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-((E)-4,7-dimethyloct-5-enyl)chrysen-2-ol (Solid **AC**): Molecular formulae  $C_{31}H_{52}O$ , white crystalline solid. Yield: (0.2337 g) 91.2%. M.pt: 131<sup>o</sup>C-132<sup>o</sup>C. Rf: 0.94 (EtOAc-Hexane, 1:4). UV [EtOH] nm (log  $\epsilon$ ): 210.79 (2.4780). IR (KBr)  $V_{max}$   $cm^{-1}$ : 3432.00 (O-H), 2939.13 (C-H), 1760.78-1641.17 (C=C), 1250.42, 1058.45 and 573.00 (fingerprint region, C-H aliphatic, C-O, C-C); <sup>1</sup>H NMR (200 MHz;  $CDCl_3$ ):  $\delta$  5.4 (1H, O-H, s), 5.2 (1H, m, ring C-H), 3.5 (1H, m, C=H alkene), 2.3 (1H, m, C=H alkene) and 0.9-1.9 (48H, m,  $CH_2$  and  $CH_3$ ). <sup>13</sup>C-NMR (50 MHz;  $CDCl_3$ ):  $\delta$  140.743 (C-5, unsaturated  $sp^2$  ring system), 121.707 (C-6, unsaturated  $sp^2$  ring system), 129.234 (C-28, unsaturated  $sp^2$ , alkene), 138.327 (C-29, unsaturated  $sp^2$ , alkene), 71.776, (C-3, C-O), 56.841 (C-14), 56.738 (C-9), 56.021 (C-4), 55.918 (C-8), 51.232 (C-13), 50.105 (C-18), 45.800 (C-7), 42.301 (C-26), 42.271 (C-25), 42.198 (C-12), 40.514 (C-10), 39.753 (C-30), 39.665 (C-23), 37.234 (C-2), 36.487 (C-17), 36.136 (C-1), 33.910 (C-24), 31.875 (C-15), 31.626 (C-16), 29.108 (C-31), 28.932 (C-32), 28.244 (C-11), 26.018 (C-27), 25.418 (C-19), 24.363 (C-20), 24.290 (C-21), 23.046 (C-22).

### Brine shrimp toxicity test

AC was subjected to brine shrimp toxicity test and  $LC_{50}$  of 1581233000.0  $\mu g/ml$  was obtained indicating non-toxicity (Table 1).

### Antioxidant activity

#### Scavenging effect on DPPH

The reduction in absorbance of DPPH at 517 nm caused by the Solid AC was measured in triplicate after 10 min. Absorbance values decreases as the concentration decreases (Table 2). Solid AC was significantly active as a free radical scavenger at 0.0625 mg/ml (64.73 %) when compared with controls, ascorbic acid, rutin and  $\alpha$ -tocopherol but lower than the activity of butylated hydroxyanisole (BHA). The percentage inhibition increases as the concen-

tration decreases unlike in the controls. DPPH is known to be a stable free radical and accepts an electron or hydrogen radical to become a stable diamagnetic molecule.

#### Scavenging effects on hydrogen peroxide ( $H_2O_2$ )

Absorbance measurement in the scavenging activity of solid AC and antioxidants standards, ascorbic acid, butylated hydroxyanisole (BHA) and  $\alpha$ -tocopherol on  $H_2O_2$  is shown in Table 3. Scavenging effect on  $H_2O_2$  was measured in triplicates after 10 min of incubation at 285 nm. At concentration of 1.0 - 0.065 mg/ml, AC scavenged hydroxyl radical in a concentration dependent manner better than the standards. Absorbance values decreases as the concentration decreases (Table 3), indicating better activity. Hydroxyl radicals are usually in-

volved in radical chain or oxidation reactions which cause diseases such as cancer, neuro-degeneration, Parkinson's disease amongst others. At 0.0625 mg/ml, AC had 99.22 % inhibition which is better than that of ascorbic acid (22.43 %), butylated hydroxyanisole (BHA) (97.0 %), and  $\alpha$ -tocopherol (86.9 %).

*Antioxidant activity by Ferric thiocyanate method (FTC)*

The amount of peroxide which oxidized ferrous chloride (FeCl<sub>2</sub>) to a reddish ferric chloride (FeCl<sub>3</sub>) pigment was determined. Here absorbance values increases as the

concentration decreases. Also, the concentration of peroxide decreases as the antioxidant activity increases. At 1.0- 0.0625 mg/ml, Solid AC showed antioxidant activities in a concentration dependent manner (Table 4). And activity was significant and better (65.23 %) at 1.0 mg/ml than the activity of BHA (17.95 %).

*Antimicrobial screening of solid AC*

The activity of solid AC against bacteria and fungi strains was tested at various concentrations. The zones of inhibition were measured and reported in Table 5.

**Table 1:** Result of brine shrimp toxicity of solid AC

| CONC.  | 1000 µg/ml |      | 100 µg/ml |      | 10 µg/ml |      | LC <sub>50</sub> µg/ml |
|--------|------------|------|-----------|------|----------|------|------------------------|
| Sample | Survivor   | Dead | Survivor  | Dead | Survivor | Dead |                        |
| AC     | 25         | 5    | 26        | 4    | 27       | 3    | 1581233000.0           |

LC<sub>50</sub> < 1000 µg/ml = Toxic, LC<sub>50</sub> > 1000 µg/ml = Non-Toxic

**Table 2:** Free radical scavenging effect of solid AC on DPPH at 517 nm\*

| CONC (mg/ml) | SOLID AC      | ASCORBIC ACID  | BHA            | ALPHA TOCOPHEROL | RUTIN         |
|--------------|---------------|----------------|----------------|------------------|---------------|
| 1.0          | 0.584 ± 0.015 | 0.0843 ± 0.010 | 0.0370 ± 0.006 | 0.6800 ± 0.029   | 0.085 ± 0.015 |
| 0.5          | 0.530 ± 0.004 | 0.2893 ± 0.128 | 0.0460 ± 0.006 | 0.7040 ± 0.003   | 0.146 ± 0.025 |
| 0.25         | 0.508 ± 0.006 | 0.2977 ± 0.124 | 0.0483 ± 0.002 | 0.7047 ± 0.007   | 0.184 ± 0.029 |
| 0.125        | 0.340 ± 0.003 | 0.32 ± 0.08200 | 0.0490 ± 0.004 | 0.7070 ± 0.007   | 0.552 ± 0.123 |
| 0.0625       | 0.329 ± 0.003 | 0.5147 ± 0.015 | 0.0650 ± 0.003 | 0.7207 ± 0.012   | 0.591 ± 0.035 |

CONC - Concentration, solid AC - an isolate of *Laportea aestuans*, DPPH - 2, 2-diphenyl-1-picrylhydrazyl radical, BHA - Butylated hydroxyl anisole

**Table 3:** Free radical scavenging effect of solid AC on H<sub>2</sub>O<sub>2</sub> at 285 nm\*

| CONC (mg/ml) | SOLID AC        | ASCORBIC ACID  | BHA            | ALPHA TOCOPHEROL |
|--------------|-----------------|----------------|----------------|------------------|
| 1.0          | 1.3392 ± 0.0007 | 0.1952 ± 0.001 | 0.0413 ± 0.016 | 0.0321 ± 0.045   |
| 0.5          | 0.8048 ± 0.0005 | 0.2078 ± 0.012 | 0.0617 ± 0.019 | 0.0633 ± 0.032   |
| 0.25         | 0.2557 ± 0.0050 | 1.2645 ± 0.119 | 0.0740 ± 0.015 | 0.1552 ± 0.061   |
| 0.125        | 0.1061 ± 0.0011 | 2.7586 ± 0.049 | 0.0947 ± 0.003 | 0.1807 ± 0.015   |
| 0.0625       | 0.0291 ± 0.0007 | 2.9236 ± 0.211 | 0.1126 ± 0.014 | 0.4940 ± 0.017   |

CONC - concentration, solid AC - an isolate of *Laportea aestuans*, BHA - Butylated hydroxyl anisole



**Table 4:** Free radical scavenging effect of solid AC on Ferric thiocyanate at 500 nm\*

| CONC (mg/ml) | SOLID AC      | ASCORBIC ACID | BHA           | ALPHA TOCOPHEROL |
|--------------|---------------|---------------|---------------|------------------|
| 1.0          | 0.261 ± 0.006 | 0.275 ± 0.006 | 0.616 ± 0.005 | 0.195 ± 0.023    |
| 0.5          | 0.311 ± 0.005 | 0.287 ± 0.050 | 0.647 ± 0.004 | 0.294 ± 0.004    |
| 0.25         | 0.324 ± 0.027 | 0.367 ± 0.004 | 0.653 ± 0.008 | 0.340 ± 0.069    |
| 0.125        | 0.333 ± 0.007 | 0.516 ± 0.008 | 0.747 ± 0.003 | 0.360 ± 0.005    |
| 0.0625       | 0.375 ± 0.016 | 0.668 ± 0.002 | 0.750 ± 0.001 | 0.377 ± 0.008    |

CONC - concentration, solid AC - an isolate of *Laportea aestuans*, BHA - Butylated hydroxyl anisole

**Table 5:** Antimicrobial screening of solid AC

| Conc | Kleb | E.coli | S. au | B.sub | S.typ | Ps.a | C.alb | A.flav | F. mon |
|------|------|--------|-------|-------|-------|------|-------|--------|--------|
| 1    | 9.5  | 11.0   | 12.0  | 14.0  | 14.0  | 10.0 | 12.5  | 10.5   | 10.0   |
| 2    | 10.5 | 10.0   | 10.0  | 10.0  | 11.0  | 9.0  | 8.5   | 10.5   | -      |
| 3    | 8.0  | 10.0   | -     | 9.0   | 9.0   | 7.0  | -     | 10.0   | -      |
| 4    | -    | 8.0    | -     | -     | 8.0   | -    | -     | -      | -      |
| Gen  | 22.0 | 23.0   | 21.5  | 21.0  | 26.0  | -    | -     | -      | -      |
| Nyst | -    | -      | -     | -     | -     | -    | 10.0  | 31.0   | 19.0   |
| Ea   | -    | -      | -     | -     | -     | 10.0 | 7.5   | -      | -      |

\*Integers 1 – 4 represent the concentrations of 1000 µg/ml, (2) 500 µg/ml, (3) 250 µg/ml, (4) 125 mg/ml. The negative control is Ea- ethyl acetate (80 µg/ml) while the positive control is gen – gentamycin (80 µg/ml) for bacteria and nyst – nystatin (80 µg/ml) for fungi. “-” represents no inhibition.

*S. au* = *staphylococcus aureus*, *E. coli* = *Escherichia coli*, *B. sub.* = *Bacillus subtilis*, *Ps.a* = *Pseudomonas aeruginosa*, *Kleb* = *Klebsiellae pneumoniae*, *S. typ* = *Salmonellae typhi*, *C.alb* = *Candida albicans*, *A.flav*– *A. flavus* and *F. mon*– *Fusarium moniliformes*

## DISCUSSION

The UV absorption spectra measured between wavelengths 190 nm and 400 nm showed the presence of chromophore in solid AB at  $\lambda_{\max}$  207.78 nm and 254.48 nm. The IR spectrum of solid AB exhibits absorption at 3432  $\text{cm}^{-1}$ , a broad band and intense peak which corresponds to an O-H group which is hydrogen bonding. The C-H stretch of methyl group absorbed at 2908.43  $\text{cm}^{-1}$ . The absorption at 1635.07  $\text{cm}^{-1}$  corresponds to C = C stretch. The several band observed within the range of 1402 - 568  $\text{cm}^{-1}$  were due to C-H aliphatic out of plane bending, C - O, C - C bonds stretching vibrations in the fingerprint region. The  $^1\text{H}$  NMR spectrum showed a methyl proton signal which resonated at  $\delta$  0.92 ppm and hydroxylic proton at  $\delta$  3.75 ppm corresponding to hydroxyl carbon at  $\delta$  76.374 ppm. The excessively tall singlet signal at  $\delta$  1.4 ppm implies that many

$\text{CH}_3$  of the same chemical environment are present which corresponds to peaks at  $\delta$  14.143 - 29.371 ppm in  $^{13}\text{C}$  NMR and 63.122 ppm corresponds to C-H of alkene. It was therefore concluded based on information obtained from  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectroscopy that the compound was (4E)-3,6-dimethylhep-4-en-3-ol.

The UV spectrum of AC showed absorption at  $\lambda$  210.79 nm. The IR ( $\text{KBr cm}^{-1}$ ) of crystal AC showed a broad band and intense peak at 3432  $\text{cm}^{-1}$  which corresponds to an O - H group which is hydrogen bonding. The C - H stretch of methyl group absorbed at 2939.13  $\text{cm}^{-1}$ . The absorption at 1657.03  $\text{cm}^{-1}$  corresponds to C = C stretch. There is also absorption at 1250.42  $\text{cm}^{-1}$  that indicated a C - O stretching vibration. The peak at 1371.28  $\text{cm}^{-1}$  corresponds to C - H deformation of  $\text{sp}^3$ . Characterization and composition of chrysen-2-ol is therefore supported by IR spectroscopy data of

hydroxyl band at  $3432.00\text{ cm}^{-1}$  typical of a chrysen-2-ol nucleus and the broad peaks representing the free 6-methyl observed on the derivative at  $2939.13\text{ cm}^{-1}$ . The NMR spectrum was in accordance with the proposed structure. The OH signal occurring at  $\delta\ 5.4$  is a singlet. The  $^{13}\text{C}$  spectrum revealed two signals characteristic of an olefinic carbon which is quaternary carbon at  $\delta\ 140.39$  and  $138.334$ , a protonated carbon. Oxygenated carbon at  $\delta\ 71.776$  indicated the presence of one hydroxyl group and the signals between  $\delta\ 23$  and  $24$  ppm are the angular methyl carbon. The two methine signals at  $\delta\ 129.241$  and  $\delta\ 121.715$  ppm, correlating with proton signal at  $\delta\ 5.2$  indicate the presence of double bond at ring. The signals between  $\delta\ 23.046$  and  $56.745$  correspond with the several proton peaks between  $\delta\ 0.9$  and  $1.9$  which indicate the presence of carbon and proton of methine and methylene group of the cyclic ring. The compound is suspected to be

1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-((E)-4,7-dimethyloct-5-enyl)chrysen-2-ol (AC). The UV, IR and NMR spectra of the compounds support similar compounds in literature. Based on the observed spectra characteristic of the two compounds, it was suspected that

4E) - 3, 6-dimethylhep-4-en-3-ol is a cleavage product of

1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-((E)-4,7-dimethyloct-5-enyl)chrysen-2-ol

The result of the brine shrimp lethality test showed that solid AC with  $\text{LC}_{50}$  values  $1581233000.0\ \mu\text{g/ml}$  was not toxic as it had  $\text{LC}_{50}$  value greater than  $1000\ \mu\text{g/ml}$ . *In vitro* antioxidant screening by the three complementary tests, namely DPPH free radical scavenging effect, scavenging effect on hydrogen peroxide and scavenging effect on ferric thiocyanate of AC showed an overall activity comparable to the antioxidant standards. It was observed that the % inhibition increased with decrease in concentration during the DPPH scavenging activity.

At  $1.0\text{ mg/ml}$ , the % inhibition was  $37.46\%$  while that of the lowest concentration ( $0.0625\text{ mg/ml}$ ) was  $64.73\%$ . It was also observed that the AC had better inhibition when compared to  $\alpha$ -tocopherol.

At  $0.125\text{ mg/ml}$  and  $0.0625\text{ mg/ml}$ , solid AC also showed better activity than rutin. The scavenging effect of solid AC on hydrogen peroxide showed that the % inhibition increased with decrease in concentration. At the highest concentration ( $1.0\text{ mg/ml}$ ), AC had % inhibition of  $64.47\%$  on hydrogen peroxide and  $99.23\%$  at the least concentration ( $0.0625\text{ mg/ml}$ ). Solid AC showed better inhibition than ascorbic acid at  $0.25\text{ mg/ml}$ ,  $0.125\text{ mg/ml}$  and  $0.0625\text{ mg/ml}$  and more inhibitive effect than butylated hydroxyl anisole (BHA) and  $\alpha$ -tocopherol at  $0.125\text{ mg/ml}$  and  $0.0625\text{ mg/ml}$ . The result of the scavenging effect of solid AC on ferric thiocyanate showed that the % inhibition increased with increase in concentration with the highest concentration  $1.0\text{ mg/ml}$  having a % inhibition of  $65.23\%$  and the least concentration which was  $0.0625\text{ mg/ml}$  having a % inhibition of  $50.00$ . It showed higher inhibition than butylated hydroxyl anisole (BHA), ascorbic acid and  $\alpha$ -tocopherol at  $0.25 - 0.0625\text{ mg/ml}$ . Activity observed was linked to the presence of chrysen-2-ol nucleus.

Antimicrobial screening result showed that solid AC was active on all the tested organisms at  $1000\ \mu\text{g/ml}$ , but as the concentration decreases the diameter of inhibition decreases. AC had better inhibition on *C. albicans* ( $12.5\text{ mm}$ ) than nystatin ( $10.0\text{ mm}$ ) indicating a selective inhibition on fungal strains.

## CONCLUSION

*Laportea aestuans* is an herb used in Africa to treat diseases related to oxidative stress and microbial infections locally. Two compounds

(4E)-3,6-dimethylhep-4-en-3-ol (AB) and 1,2,3,4,4a,4b,5,6,6a,7,8,9,10,10a,10b,11-hexadecahydro-1,1,6a,10b-tetramethyl-7-

((*E*)-4,7-dimethyloct-5-enyl)chrysen-2-ol (AC) were isolated and characterized using data obtained from UV/Visible, IR, <sup>1</sup>H and <sup>13</sup>C NMR spectroscopies. AC was found to be non-toxic and possessed significant antioxidant activity when compared to standards; ascorbic acid, butylated hydroxyl anisole, rutin and  $\alpha$ -tocopherol. The selective antimicrobial activity of AC confirmed its usefulness in the treatment of various infectious diseases caused by bacteria and fungi locally.

### ACKNOWLEDGEMENT

The authors gladly acknowledge the staff of Multi Research Central Science Laboratory, University of Ibadan, Oyo State and Central Science Laboratory Obafemi Awolowo University (OAU) Ile Ife, Osun State for the use of spectroscopic equipment. The authors will also like to thank Dr Idowu of Department of pharmaceutical Chemistry, OAU, Ile Ife and staff of the department of Microbiology University of Ibadan, Nigeria for assisting in carrying out the antimicrobial analysis.

### REFERENCES

Abu-Rabia A. Herbs as a food and medicine source in Palestine. *Asian Pac J Cancer Prev* 2005;6:404-7.

Alan L, Miller ND. Antioxidant flavonoids: structure, function and clinical usage. *Alt Med Rev* 1996;1:103-13.

Ames BN, Shigenaga MK, Hagen TM. Oxidants, antioxidants, and the degenerative diseases of aging. *Proc Natl Acad Sci USA* 1993;90:7915-22.

Bayer AW, Kirby MDK, Sheris JC, Trick M. Antibiotic susceptibility testing by single disc diffusion method. *J Clin Pathol* 1986;45:493-6.

Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* 2007;297:842-57. Erratum in: *JAMA* 2008; 299:765-6.

Burkill HM. The useful plants of west tropical Africa (2<sup>nd</sup> ed.). Vol. 1: Royal Botanical Garden. UK, KEW 1985:641-3.

Cadenas E, Davies KJA. Mitochondrial free radical generation, oxidative stress, and aging. *Free Rad Biol Med* 2000;29:222-30.

Chew WL. A monograph of *Laportea* (Urticaceae). *Gard Bull Straits Settlements* 1969;25: 111-78.

de la Fuente M, Victor M. Antioxidants as modulators of immune function. *Immunol Cell Biol* 2000;78:49-54.

Di Matteo V, Esposito E. Biochemical and therapeutic effects of antioxidants in the treatment of Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. *Curr Drug Target CNS Neurol Disord* 2003;2:95-107.

Dokosi OB. Herbs of Ghana. Accra, Ghana: Council for Industrial Research, 1998 (pp 510-1).

Duraipandiyar V, Ayyanar M, Ignacimuthu S. Antimicrobial activity of some ethnomedicinal plants used by paliyar tribe from Tamil Nadu, India. *BMC Comp Alt Med* 2006;6:35-41.

Falope MO, Ibrahim H, Takeda Y. Screening of higher plants requested as pesticides using the brine shrimp lethality assay. *Int J Pharmacognosy* 1993;37:230-54.

Brinker FJ. Herb contraindications and drug interactions (2<sup>nd</sup> ed.). Sandy, Oregon: Eclectic Medical Publ., 1998.

Friis I. Urticaceae. In: Kubitzki K et al. (eds.): The families and genera of vascular plants, Vol. 2: Flowering plants, dicotyledons: magnoliid, hamamelid and caryophyllid families: dicotyledons (pp 612-30). Berlin: Springer-Verlag, 1993.

Gbile ZO. Ethnobotany, taxonomy and conservation of medicinal plants. In: Sofowora A (ed.): The state of medicinal plants research in Nigeria (pp 267-9). Ibadan, Nigeria: University of Ibadan Press, 1986.

Ghani AK. Medicinal plants and traditional medicine portions: problems and prospects of their standardization. In: Sofowora A (ed.): The state of medicinal plants research in Nigeria (pp 65-77). Ibadan, Nigeria: University of Ibadan Press, 1986.

Gilbert JC, Martin SF. Experimental organic chemistry: a miniscale and microscale approach (4<sup>th</sup> ed.). Belmont, CA: Brooks/Cole, Cengage Learning, 2006 (pp 171-90).

Gulcin IM, Oktay OK, Aslan A. Determination of antioxidant activity of *Linchen Cetraria islandica* (L). Ac. J Ethnopharmacol 2002;79:325-9.

Hatano T, Kagawa H, Yasuhora T, Okuta T. Two new flavonoids and other constituents in licorice root: Their relative astringency and radical scavenging effects. Chem Pharm Bull 1988;36:2090-7.

Heywood VH. Flowering plants of the world. New York: Oxford Univ. Press, 1993.

Keddy CJ, Green JC, Bonnel MA. Review of whole organism bioassays: soil, water and freshwater assessment in Canada. Ecotoxicol Environ Saf 1995;30:221-51.

Knight JA. Free radicals: their history and current status in aging and disease. Ann Clin Lab Sci 1998;28:331-46.

Kokwaro JO. Medicinal plants of East Africa. Kenya: Kenya Literature Bureau, 1993.

Lans CA. Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus. J Ethnobiol Ethnomed 2006;2:45.

Lans CA. Ethnomedicines used in Trinidad and Tobago for reproductive problems. J Ethnobiol Ethnomed 2007;3:13.

Meyer BN, Ferrign RN, Putnam JE, Jacobson LB, Nicholas DE, McLaughlin JL. Brine shrimp: a convenient general bioassay for active plant constituents. Planta Med 1982;45:31-4.

Miller RA, Britigan BE. Role of oxidants in microbial pathophysiology. Clin Microbiol Rev 1997;10:1-18.

Morrison JF, Twumasi SK. Comparative studies on the in vitro antioxidant properties of methanolic and hydro-ethanolic leafy extracts from eight edible leafy vegetables of Ghana. Afr J Biotech 2010;9:5177-84.

Mutee AF, Salhimi SM, Yam MF, Lim CP, Abdullah GZ, Ameer OZ et al. In vivo anti-inflammatory and in vitro antioxidant activities of *Peperomia pellucida*. Int J Pharmacol 2010;6:686-90.

Nadine A. General information for medicinal plants. New edition Vol. 6 West India Chew Western edition, 2001;5: 7-31.

Oloyede GK. Toxicity, antimicrobial and antioxidant activities of methyl salicylate dominated essential oils of *Laportea aestuans* (Gaud). Arab J Chem 2011;(in press). <http://dx.doi.org/10.1016/j.arabjc.2011.09.019>

Oloyede GK, Farombi OE. Antioxidant properties of *crinum ornatum* Bulb Extract. World J Chem 2010;5:32-6.

Oloyede GK, Oke MJ, Raji Y, Olugbade AT. Antioxidant and anticonvulsant alkaloids in *Crinum ornatum* Bulb Extract. *World J Chem* 2010;5:26-31.

Oloyede GK, Ayanbadejo OE. Phytochemical, toxicity, antimicrobial and antioxidant screening of extracts obtained from *Laportea aestuans* (Gaud). *J Med Sci (Science Alert)* 2013;(in print).

Paul J. Constituent *Laportea aestuans* and *Serjania Paucidentata*. Thesis. Saint Augustine, Trinidad and Tobago: University of the West Indies, 2008.  
<http://hdl.handle.net/2139/2238>.

Pinzino C, Capocchi A, Galleschi L, Saviozzi F, Nanni B, Zandomenighi M. Aging, free radicals, and antioxidants in wheat seeds. *J Agric Food Chem* 1999;47:1333-9.

Pourmorad F, Hosseinimehr SJ, Shahabimajd N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol* 2006;5:1142-5.

Schmidt EJ, Boswell JS, Walsh JP, Schellenberg MM, Winter TE, Li C et al. Activities of cholic acid-derived antimicrobial agents against multidrug resistant bacteria. *J Antimicrob Chemother* 2006;47:671-4.

Sies H. Oxidative stress: oxidants and antioxidants. *Exp Physiol* 1997;82:291-5.

Soares JR, Dinis TC, Cohn AP, Almeida LM. Antioxidant activity of some extracts of *Thymus zygis*. *Free Radic Res* 1997;6:469-78.

Stadtman ER. Protein oxidation and aging. *Science* 1992;257:1220-4.

Stanner SA, Hughes J, Kelly CN, Buttriss J. A review of the epidemiological evidence for the antioxidant hypothesis. *Public Health Nutr* 2004;7:407-22.