

Guest editorial:

THE REDISCOVERY OF HepG2 CELLS FOR PREDICTION OF DRUG INDUCED LIVER INJURY (DILI)

Ahmed Ghallab

Department of Forensic and Toxicology, Faculty of Veterinary Medicine,
South Valley University, Qena, Egypt; ghallab@vet.svu.edu.eg

In the past HepG2 cells have been considered to show only little similarity to primary human hepatocytes (Godoy et al., 2013; Hewitt et al., 2007). However, recently it has become clear that the phenotype of HepG2 cells strongly depends on culture conditions. In a recent study, Ramaiahgari et al. (2014) cultivated HepG2 cells in spheroids. In this culture system the cells stopped proliferation and strongly upregulated phase I and phase II drug metabolizing enzymes and transporters (Ramaiahgari et al., 2014). Moreover, albumin and urea metabolizing enzymes were upregulated. This study shows that the potential of HepG2 cells may have been underestimated in the past. A critical aspect seems to be that they have to be kept in three dimensional culture systems.

Much effort has been invested in the establishment of hepatocyte culture systems that help to identify hepatotoxic compounds (Ilkavets, 2013; Abdelhamid et al., 2013; Vinken et al., 2013; Hasmall et al., 2001; Waterfield et al., 1998; Mennes et al., 1994; Krijt et al., 1993; Godoy et al., 2009; Adler et al., 2014; Maruf and O'Brien, 2014). In recent years HepG2 cells have become more and more popular for this purpose (Mostafavi-Pour et al., 2013; Shan et al., 2013; Krithika et al. 2013; Doricakova et al., 2013; Straser et al. 2011; Dias da Silva et al., 2013; Horinouchi et al., 2014; Wang et al., 2014). One reason for their frequent application may be that they are freely available in contrast to some other cell lines obtained from

human liver tumors that are only commercially available. Another successful application of HepG2 cells is expression of human genes (Lahoz et al., 2013). The group of Castell and Gomez-Lechon have specialized in this field (Tolosa et al., 2012; Donato et al., 2010; 2008). Recently, five P450 enzymes have been expressed simultaneously in HepG2 cells resulting in a powerful test system for compounds that require metabolic activation (Tolosa et al., 2013). It is clear that primary human hepatocytes currently still represent the gold standard for hepatotoxicity testing (Hengstler et al., 2000; 2009; Hewitt et al., 2007). Future studies will have to show to which degree HepG2 based culture system can be used to predict hepatotoxic compounds in vitro.

REFERENCES

- Abdelhamid G, Amara IE, Anwar-Mohamed A, El-Kadi AO. Modulation of cytochrome P450 1 (Cyp1) by vanadium in hepatic tissue and isolated hepatocyte of C57BL/6 mice. *Arch Toxicol.* 2013;87:1531-43.
- Adler M, Leich E, Ellinger-Ziegelbauer H, Hewitt P, Dekant W, Rosenwald A, et al. Application of RNA interference to improve mechanistic understanding of omics responses to a hepatotoxic drug in primary rat hepatocytes. *Toxicology.* 2014;326C:86-95.
- Dias da Silva D, Carmo H, Lynch A, Silva E. An insight into the hepatocellular death induced by amphetamines, individually and in combination: the involvement of necrosis and apoptosis. *Arch Toxicol.* 2013;87:2165-85.

- Donato MT, Lahoz A, Castell JV, Gómez-Lechón MJ. Cell lines: a tool for in vitro drug metabolism studies. *Curr Drug Metab.* 2008;9:1-11.
- Donato MT, Hallifax D, Picazo L, Castell JV, Houston JB, Gomez-Lechón MJ, et al. Metabolite formation kinetics and intrinsic clearance of phenacetin, tolbutamide, alprazolam, and midazolam in adenoviral cytochrome P450-transfected HepG2 cells and comparison with hepatocytes and in vivo. *Drug Metab Dispos.* 2010;38:1449-55.
- Doricakova A, Novotna A, Vrzal R, Pavek P, Dvorak Z. The role of residues T248, Y249 and T422 in the function of human pregnane X receptor. *Arch Toxicol.* 2013;87:291-301.
- Godoy P, Hengstler JG, Ilkavets I, Meyer C, Bachmann A, Müller A, et al. Extracellular matrix modulates sensitivity of hepatocytes to fibroblastoid dedifferentiation and transforming growth factor beta-induced apoptosis. *Hepatology.* 2009;49:2031-43.
- Godoy P, Hewitt NJ, Albrecht U, Andersen ME, Ansari N, Bhattacharya S, et al. Recent advances in 2D and 3D in vitro systems using primary hepatocytes, alternative hepatocyte sources and non-parenchymal liver cells and their use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. *Arch Toxicol.* 2013;87:1315-530.
- Hasmall S, James N, Hedley K, Olsen K, Roberts R. Mouse hepatocyte response to peroxisome proliferators: dependency on hepatic nonparenchymal cells and peroxisome proliferator activated receptor alpha (PPARalpha). *Arch Toxicol.* 2001;75:357-61.
- Hengstler JG1, Utesch D, Steinberg P, Platt KL, Diener B, Ringel M, et al. Cryopreserved primary hepatocytes as a constantly available in vitro model for the evaluation of human and animal drug metabolism and enzyme induction. *Drug Metab Rev.* 2000;32:81-118.
- Hengstler JG, Godoy P, Bolt HM. The dilemma of cultivated hepatocytes. *Arch Toxicol.* 2009;83:101-3.
- Hewitt NJ, Lechón MJ, Houston JB, Hallifax D, Brown HS, Maurel P, et al. Primary hepatocytes: current understanding of the regulation of metabolic enzymes and transporter proteins, and pharmaceutical practice for the use of hepatocytes in metabolism, enzyme induction, transporter, clearance, and hepatotoxicity studies. *Drug Metab Rev.* 2007;39:159-234.
- Horinouchi Y, Summers FA, Ehrenshaft M, Mason RP. Free radical generation from an aniline derivative in HepG2 cells: A possible captodative effect. *Free Radic Biol Med.* 2014;78C:111-7.
- Ilkavets I. A special issue about hepatotoxicity and hepatocyte in vitro systems. *Arch Toxicol.* 2013;87:1313-4.
- Krijt J, van Holsteijn I, Hassing I, Vokurka M, Blaauboer BJ. Effect of diphenyl ether herbicides and oxadiazon on porphyrin biosynthesis in mouse liver, rat primary hepatocyte culture and HepG2 cells. *Arch Toxicol.* 1993;67:255-61.
- Krithika R, Verma RJ, Shrivastav PS. Antioxidative and cytoprotective effects of andrographolide against CCl4-induced hepatotoxicity in HepG2 cells. *Hum Exp Toxicol.* 2013;32:530-43.
- Lahoz A, Vilà MR, Fabre M, Miquel JM, Rivas M, Maines J, et al. An in vitro tool to assess cytochrome P450 drug biotransformation-dependent cytotoxicity in engineered HepG2 cells generated by using adenoviral vectors. *Toxicol In Vitro.* 2013;27:1410-5.
- Maruf AA, O'Brien P. Flutamide-induced cytotoxicity and oxidative stress in an in vitro rat hepatocyte system. *Oxid Med Cell Longev.* 2014;2014:398285.
- Mennes WC, Wortelboer HM, Hassing GA, van Sandwijk K, Timmerman A, Schmid BP, et al. Effects of clofibrate and beclobric acid in rat and monkey hepatocyte primary culture: influence on peroxisomal and mitochondrial beta-oxidation and the activity of catalase, glutathione S-transferase and glutathione peroxidase. *Arch Toxicol.* 1994;68:506-11.
- Mostafavi-Pour Z, Khademi F, Zal F, Sardarian AR, Amini F. In vitro analysis of CsA-induced hepatotoxicity in HepG2 cell line: oxidative stress and $\alpha 2$ and $\beta 1$ integrin subunits expression. *Hepat Mon.* 2013;13(8):e11447.
- Ramaiahgari SC1, den Braver MW, Herpers B, Terpstra V, Commandeur JN, van de Water B, et al. A 3D in vitro model of differentiated HepG2 cell spheroids with improved liver-like properties for repeated dose high-throughput toxicity studies. *Arch Toxicol.* 2014;88:1083-95.
- Shan G, Ye M, Zhu B, Zhu L. Enhanced cytotoxicity of pentachlorophenol by perfluorooctane sulfonate or perfluorooctanoic acid in HepG2 cells. *Chemosphere.* 2013;93:2101-7.
- Straser A, Filipič M, Zegura B. Genotoxic effects of the cyanobacterial hepatotoxin cylindrospermopsin in the HepG2 cell line. *Arch Toxicol.* 2011;85:1617-26.
- Tolosa L, Donato MT, Pérez-Cataldo G, Castell JV, Gómez-Lechón MJ. Upgrading cytochrome P450 activity in HepG2 cells co-transfected with adenoviral vectors for drug hepatotoxicity assessment. *Toxicol In Vitro.* 2012;26:1272-7.

Tolosa L, Gómez-Lechón MJ, Pérez-Cataldo G, Castell JV, Donato MT. HepG2 cells simultaneously expressing five P450 enzymes for the screening of hepatotoxicity: identification of bioactivable drugs and the potential mechanism of toxicity involved. *Arch Toxicol.* 2013;87:1115-27.

Vinken M, Maes M, Cavill R, Valkenburg D, Ellis JK, Decrock E, et al. Proteomic and metabolomic responses to connexin43 silencing in primary hepatocyte cultures. *Arch Toxicol.* 2013;87:883-94.

Wang Y, Wu J, Lin B, Li X, Zhang H, Ding H, et al. Galangin suppresses HepG2 cell proliferation by activating the TGF- β receptor/Smad pathway. *Toxicology.* 2014;326C:9-17.

Waterfield CJ, Westmoreland C, Asker DS, Murdock JC, George E, Timbrell JA. Ethionine toxicity in vitro: the correlation of data from rat hepatocyte suspensions and monolayers with in vivo observations. *Arch Toxicol.* 1998;72:588-96.