

Guest editorial:

DISCOVERING URINARY BLADDER CANCER RISK VARIANTS: STATUS QUO AFTER ALMOST TEN YEARS OF GENOME-WIDE ASSOCIATION STUDIES

Silvia Selinski

Leibniz-Institut für Arbeitsforschung an der TU Dortmund,
Leibniz Research Centre for Working Environment and Human Factors (IfADo),
Ardeystrasse 67, 44139 Dortmund, Germany
selinski@ifado.de

<http://dx.doi.org/10.17179/excli2017-1000>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

About ten years ago Kiemeny and colleagues (2008) published the first genome-wide association study (GWAS) discovering two novel single nucleotide polymorphisms (SNPs) near *MYC* (rs9642880) and *TP63* (rs710521) associated with urinary bladder cancer (UBC) risk. Meanwhile, further GWAS and candidate gene studies identified and confirmed a number of susceptibility variants for UBC (Selinski, 2012, 2013, 2014; Dudek et al., 2013; Selinski et al., 2013; Golka et al., 2011). Currently, fifteen genomic regions seem to play a major role in development of this disease (Table 1). It can be assumed that almost all relevant polymorphisms have been discovered now. The most recent UBC GWAS of Figueroa et al. (2016) required 15,058 cases and 286,270 controls to discover a further susceptibility region at 13q34 (*MCF2L* gene), including the Icelandic and the Dutch GWAS (Gudbjartsson et al., 2015; Kiemeny et al., 2008). The odds ratios of the most recent UBC risk SNPs are rather small, e.g. the *MCF2L* intron variant rs4907479, with the strongest signal in the 2016 fine-mapping study of Figueroa et al. (2016) resulted in an odds ratio (OR) of 1.13.

Several UBC risk variants are particularly relevant in persons exposed to bladder

carcinogens – mainly by tobacco smoke (Burger et al., 2013; Garcia-Closas et al., 2005, 2011, 2013; Selinski et al., 2013; Moore et al., 2011) but also by occupation and environment (Ebbinghaus et al., 2017; Höhne et al., 2017; Krech et al., 2017; Lukas et al., 2017; Carreón et al., 2014; Golka et al., 2012, 2009, 2004, 2002, 1997, 1996; Delclos and Lerner, 2008; Ovsianikov et al., 2012; Rushton et al., 2012). However, currently no particular variant that is only relevant in exposed persons could be identified and replicated in GWAS. In 2014, Figueroa et al. (2014a) identified two variants in a genome-wide smoking × SNP interaction study –rs1711973 (*FOXF2*) relevant for never smokers and rs12216499 (*RSPH2-TAGAP-EZR*) in ever smokers, but both could not be confirmed using a large replication series (Figueroa et al., 2016). Nevertheless, interaction analyses and stratification regarding smoking habits and cancer invasiveness are promising future approaches to uncover further susceptibility variants that are relevant for particular subgroups of the UBC patients.

A further challenge is a genome-wide search for variants associated with bladder

cancer recurrence and progression. This requires a large number of UBC patients with follow-up of several years after first diagnosis. However, it can be assumed that variants relevant for UBC development are also associated with UBC recurrence. Recurrence of this tumor occurs in approximately half of the patients with a median recurrence-free time of almost one year. Selinski et al. (2017a) showed that the ultra-slow N-acetyltransferase 2 (*NAT2*) genotype was associated with a significant reduction of recurrence-free time (8.4 months) compared to rapid acetylators (11 months) and an increased recurrence risk (66 % vs. 50 %, OR=1.89, 95 % CI = 1.06–3.38). Effects were more pronounced in ultra-slow smokers (7.9 months, 73 %) indicating the relevance of gene-environment interaction also for prognosis. Correspondingly, Lukas et al. (2017) showed in a series of 143 UBC cases with suspected occupational bladder cancer the importance of co-occurring susceptibility variants, particularly co-occurring *GSTM1* negative and rs11892031[A/A] for UBC recurrence. They discovered that UBC cases with an elevated number of risk alleles had a significantly shorter median relapse-free time of 8 months compared to cases with few risk alleles.

According to the different importance of several genetic variants depending on exposure to bladder carcinogens, e.g. *GSTM1* and *NAT2*, different variant combinations seem to play a major role in smokers and never smokers (Selinski et al., 2017b; Schwender et al., 2012). Recently, Selinski et al. (2017b) identified and replicated in a large multicentric case-controls series (discovery series: 2969 cases / 3285 controls, replication series: 2080 cases / 2167 controls) four-variant combinations out of twelve well-known UBC risk variants. The highest odds ratios were found in never smokers with the best combination (rs1014971[AA] × rs1058396[AG,GG] × rs11892031[AA] × rs8102137[CC,CT]) resulting in an OR of 2.59 (95 % CI = 1.93-3.47; P = 1.87 × 10⁻¹⁰, frequency in never smoking cases: 25 %). Odds ratios of the best combinations found in smokers were clearly lower (current smokers: 1.56, former: 2.13, ever: 1.55) and different variant combinations were relevant, especially *GSTM1*, rs1058396 (*SLC14A1*) and rs11892031 (*UGT1A*) combinations in current and rs9642880 (*MYC*), rs1495741 (*NAT2*) and rs8102137 (*CCNE1*) combinations in former smokers (Selinski et al., 2017b).

Table 1: Currently confirmed polymorphisms that are associated with UBC risk, their association with bladder carcinogen exposure and prognosis (update of Selinski, 2014) according to Selinski (2014). Polymorphisms, associated genes and locations are printed in bold, risk alleles or genotypes are given in brackets.

Location Gene(s)	Key message	Reference
1p13.3 GSTM1 (<i>glutathione S-transferase mu 1</i>)	GSTM1 null was confirmed as risk factor for increased UBC risk in a GWAS (OR=1.47).	Rothman et al. (2010)
	GSTM1 null was associated with recurrence (<i>GSTM1</i> *present/null HR=1.5, <i>GSTM1</i> *null/null HR=2.0 vs. <i>GSTM1</i> *present/present) and mortality (<i>GSTM1</i> *null/null vs. <i>GSTM1</i> * present/null HR=1.9) in the Copenhagen City Heart Study (CCHS).	Nørskov et al. (2011)
	GSTM1 null × Smoking (additive interaction) was associated with increased UBC risk (OR _{SNP×Smoking} =4.69, P _{additive} =0.008)	Garcia-Closas et al. (2013)
	GSTM1 null was not associated with UBC invasiveness or prognosis.	Grotenhuis et al. (2014)
	GSTM1 null UBC risk was confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.37, P=2.306×10 ⁻⁴).	Wang et al. (2014)

Location Gene(s)	Key message	Reference
2q37.1 UGT1A (<i>UDP glucuronosyl-transferase 1 family, polypeptide A complex locus</i>)	Rs11892031[C] (<i>UGT1A</i> intron SNP) was associated with decreased UBC in a GWAS (OR=0.84). <i>UGT1A</i> proteins are involved in the metabolisms of bladder carcinogens via glucuronidation.	Rothman et al. (2010)
	Rs17863783[T] (<i>UGT1A6</i> exon variant, MAF 2 %) was associated with UBC in a fine-mapping study (OR=0.55) and explained most of the effect of rs11892031. Rs17863783 increased <i>UGT1A6.1</i> mRNA expression <i>in vitro</i> .	Tang et al. (2012)
	Rs11892031[C] was particularly associated with a lower UBC risk in persons with occupational exposure to aromatic amines and PAHs (OR=0.68 exposed persons, OR=0.83 total study group).	Selinski et al. (2012)
	Rs17863783[T] × Smoking showed an additive interaction with smoking (OR _{SNP×Smoking} =3.83, P _{additive} =8.8 × 10 ⁻⁴). Rs11892031[C] × Smoking was not significant.	Garcia-Closas et al. (2013)
	Rs11892031 and rs17863783 were not associated with UBC invasiveness or prognosis.	Grotenhuis et al. (2014)
3q26.2 TERC (<i>telomerase RNA component</i>) – ACTRT3 (<i>actin-related protein T3</i>) – MYNN (<i>myoneurin</i>) – LRRC34 (<i>leucine rich repeat containing 34</i>)	Rs10936599[C] (intergenic) was associated with increased UBC risk UBC in a GWAS (OR=1.18). The nearby genes MYNN , TERC and ACTRT3 were overexpressed in bladder tumor tissue compared to normal bladder tissue. However, rs10936599 did not modify the gene expression.	Figuroa et al. (2014b)
	Rs10936599[C] UBC risk was confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.22, P=7.179×10 ⁻⁴).	Wang et al. (2014)
3q28 TP63 (<i>tumor protein 63</i>)	Rs710521[A] (intergenic) was associated with UBC in a GWAS (OR=1.19).	Kiemeny et al. (2008)
	Rs710521[A] UBC risk (OR=1.16) was not modified by smoking or occupational exposure to bladder carcinogens.	Lehmann et al. (2010)
	Rs710521[A] × Smoking (interaction) was not significant in an interaction analysis.	Garcia-Closas et al. (2013)
	Rs710521 was not associated with UBC invasiveness or prognosis.	Grotenhuis et al. (2014)
	Rs710521[A] UBC risk could not be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.14, P=0.061).	Wang et al. (2014)
4p16.3 TACC3 (<i>transforming, acidic coiled-coil containing protein 3</i>) – FGFR3 (<i>fibroblast growth factor receptor 3</i>)	Rs798766[T] (<i>TACC3</i> intron SNP, 70 kb of FGFR3) was associated with UBC in a GWAS (OR=1.24).	Kiemeny et al. (2010)
	Rs798766[T] × Smoking (interaction) was not significant in an interaction analysis.	Garcia-Closas et al. (2013)
	Rs798766[T] was associated with NMIBC recurrence in non-smokers (P=2.7 × 10 ⁻⁵ , HR _{T/T} =2.71, HR _{C/T} =2.43) but not in ever smokers nor in the total case group (P=0.75 and P=0.12, respectively). Rs798766 was not associated with progression from NMIBC to MIBC nor with mortality in MIBC cases (P=0.28).	Grotenhuis et al. (2014)
	Rs798766[T] UBC risk was confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.24, P=0.012).	Wang et al. (2014)

Location Gene(s)	Key message	Reference
5p15.33 CLPTM1L (<i>cisplatin resistance related protein CRR9p</i>) – TERT (<i>telomerase reverse transcription</i>)	Rs401681[C] (CLPTM1L intron SNP) and the nearby synonymous rs2736098[A] (synonymous TERT exon SNP) were both associated with UBC in a GWAS (OR=1.12 and 1.16, respectively). CLPTM1L is involved in cisplatin-induced apoptosis. TERT is associated with telomere maintenance and aging. Both genes are located at a cancer susceptibility locus at 5p15.33.	Rafnar et al. (2009)
	Rs401681[C] × Smoking (interaction) was not significant in an interaction analysis.	Garcia-Closas et al. (2013)
	Rs401681 and rs2736098 were not associated with UBC invasiveness or prognosis.	Grotenhuis et al. (2014)
	Rs401681[C] and rs2736098[G] UBC risk could not be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.02 and 1.01, P=0.779 and 0.858, respectively).	Wang et al. (2014)
8p22 NAT2 (<i>N-acetyltransferase 2</i>)	Rs1495741[A/A] (intergenic, 14 kb 3' of NAT2 tagSNP), was associated with UBC in a GWAS (OR _{A/A} =1.15). NAT2 variants result in a reduced acetylation of bladder carcinogens and are an established susceptibility factor for UBC in persons exposed to bladder carcinogenic aromatic amines, e.g. smokers. Rs1495741[A/A] corresponds to the slow phenotype. Increased risks were observed in current smokers (OR=1.25).	Rothman et al. (2010)
	Rs1495741[A/A] showed a very good specificity (94 %) for the prediction of the NAT2 phenotype. However, a perfect specificity was obtained by the common 7-SNPs NAT2 genotype (1.00) and a 2-SNPs genotype (1.00) based on the established 7 NAT2 SNPs in Caucasians.	Selinski et al. (2011)
	Rs1495741[A] × Smoking interaction increased UBC risk significantly (OR _{SNP×Smoking} =2.48, P _{additive} =6.6 × 10 ⁻⁴ , P _{multiplicative} =0.029).	Garcia-Closas et al. (2013)
	Rs1495741[A/A] was associated with NMIBC recurrence (P=0.02, HR=1.29) but not progression in ever smokers (P=0.27) and reduced NMIBC progression risk in non-smokers (P=0.03, HR=0.42).	Grotenhuis et al. (2014)
	Rs1495741[A] UBC risk could be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.13, P=0.034).	Wang et al. (2014)
	Rs2294008[T] was associated with UBC in a GWAS (OR=1.15). Rs2294008[T] altered the start codon of PSCA leading to a reduced promoter activity <i>in vitro</i> . Overexpression of PSCA in prostate and bladder tumors is well-known.	Wu et al. (2009)
8q24.3 PSCA (<i>prostate stem cell antigen</i>)	Rs2978974[A] (in an alternative untranslated 1 st exon of PSCA) was detected in a fine-mapping study (OR=1.11) and showed a significant interaction effect (P=0.035) with rs2294008 (10 kb upstream rs2978974) instead of capturing the same signal.	Fu et al. (2012)
	Rs2294008[T] × Smoking interaction (additive) increased UBC risk significantly (OR _{SNP×Smoking} =2.90, P _{additive} =0.033).	Garcia-Closas et al. (2013)
	Rs2294008 and rs2978974 were not associated with NMIBC or MIBC prognosis.	Grotenhuis et al. (2014)
	Rs2294008 UBC risk could be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.21, P=0.003).	Wang et al. (2014)

Location Gene(s)	Key message	Reference
8q24.21 MYC (<i>v-myc myelocytomatosis viral oncogene homolog (avian)</i>)	Rs9642880[T] (intergenic, 30 kb MYC) was associated with UBC in a GWAS (OR=1.22).	Kiemeney et al. (2008)
	Rs9642880[T] UBC risk was less pronounced in cases with suspected exposure to bladder carcinogens (OR=1.04 and 1.11 exposed study groups) than in non-exposed persons (OR=1.36) in contrast to GSTM1 null (OR=2.43 and 1.38 in exposed study groups, OR=1.22 non-exposed persons).	Golka et al. (2009)
	Rs9642880[T] × Smoking interaction (additive) increased UBC risk significantly (OR _{SNP×Smoking} =3.49, P _{additive} =0.035).	Garcia-Closas et al. (2013)
	Rs9642880[G] was associated with NMIBC progression (P _{trend} =2.6 × 10 ⁻³ , HR _{G/G} =1.81, HR _{G/T} =1.06) but not with NMIBC recurrence (P=0.98) nor with mortality among MIBC cases (P=0.56).	Grotenhuis et al. (2014)
	Rs9642880[T] UBC risk could be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.24, P=0.010).	Wang et al. (2014)
11p15.5 LSP1 (<i>lymphocyte-specific protein 1, longer transcript</i>)	Rs907611[A] (130 bp upstream LSP1 transcription start site) was associated with UBC in a GWAS (OR=1.15). Rs907611[A] did not modify LSP1 and miRNA-4298 expression in bladder tissue. MIRNA-4298 is located in the LSP1 gene and in the same haploblock as rs907611 .	Figuroa et al. (2014b)
	Rs907611 UBC risk could not be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.12, P=0.129).	Wang et al. (2014)
13q34 MCF2L (<i>MCF.2 cell line derived transforming sequence like</i>)	Rs4907479 (MCF2L intron SNP) was associated with UBC in a GWAS (OR = 1.13). No association was found with MCF2L mRNA expression in bladder tumors. MCF2L encodes a guanine nucleotide exchange factor that is involved in the Rho/Rac signaling pathways.	Figuroa et al. (2016)
18q12.3 SLC14A1 (<i>solute carrier family 14 (urea transporter), member 1 (Kidd blood group)</i>)	Rs1058396[G] (SLC14A1 exon 8 missense SNP) and rs17674580[T] (SLC14A1 intron SNP) both increased UBC risk in a GWAS (OR=1.14, OR=1.17) but captured the same signal. Rs17674580 explained the effect of rs1058396. SLC14A1 plays a role in maintenance of a constant urea concentration gradient in the kidney. The exon missense SNP rs1058396 (D280N) determines two alleles of the Kidd blood system.	Rafnar et al. (2011)
	Rs7238033[T] (OR=1.20) as well as rs10775480[T] and rs10853535[C] (both: OR=1.16) were associated with UBC in a GWAS. Rs1058396 was in strong LD with rs7238033 , rs10775480 and rs11082469 (r ² =0.71, 0.64, and 0.93, respectively). The highly correlated rs10775480[T] and rs10853535[C] (r ² =1.00) were used as mutual tagSNPs.	Garcia-Closas et al. (2011)
	rs10775480[T]/rs10853535[C] × Smoking interactions were not significant in an interaction analysis (P _{additive} =0.053, P _{multiplicative} =0.833).	Garcia-Closas et al. (2013)
	Rs1058396 was not associated with UBC prognosis.	Grotenhuis et al. (2014)
	Rs17674580[C] UBC risk could be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.65, P=8.507×10 ⁻⁸).	Wang et al. (2014)

Location Gene(s)	Key message	Reference
19q12 CCNE1 (<i>cyclin E1</i>)	Rs8102137[C] (intergenic) transition was associated with UBC in a GWAS (OR=1.13). CCNE1 is involved in the cell cycle G1/S phase CCNE1 over-expression seemed to be associated with tumorigenesis and UBC prognosis.	Rothman et al. (2010)
	Rs8102137[C] × Smoking did not modify UBC risk in an interaction analysis ($P_{\text{additive}}=0.961$, $P_{\text{multiplicative}}=0.133$).	Garcia-Closas et al. (2013)
	Rs8102137 was not associated with UBC prognosis.	Grotenhuis et al. (2014)
	Rs8102137[C] UBC risk could not be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.12, $P=0.253$).	Wang et al. (2014)
20p12.2 JAG1 (<i>jagged 1</i>)	Rs6104690[A] (intergenic, 300 kb of JAG1) was associated with UBC in a GWAS (OR=1.12).	Figuroa et al. (2014b)
	Rs62185668[A] and rs4813953[T] (OR=1.19 and 1.16, resp.) as well as the indel rs148953085 (all: intergenic, 300 kb of JAG1) were associated with UBC in a GWAS. Rs62185668 represented the signal best. Rs62185668 and rs148953085 (both in a predicted DNaseI hotspot) showed a high correlation ($r^2=0.96$). The correlation between rs62185668 and rs4813953 was moderate ($r^2=0.50$). The effect of rs6104690[A] could be fully explained by the nearby SNP rs62185668 (rs6104690 : $OR_{\text{adj}}=0.99$, $P_{\text{adj}}=0.85$, $r^2=0.30$). Expression of the NOTCH ligand JAG1 was reduced in bladder tumors compared to normal urothelium. Rs62185668[A] led to a reduced expression of JAG1 in low-passage urothelial cells.	Rafnar et al. (2014)
	Rs6108803 (intergenic, 300 kb of JAG1) was associated with UBC in a fine-mapping study (OR=1.18) and explained most of the effect of rs62185668 ($P_{\text{adj}}=0.25$). The effect of rs6104690 (OR=1.11) could be confirmed in 8,147 additional cases and 274,456 controls (in total: 15,058 / 286,270) compared to Figuroa et al. (2014b). Rs6108803-rs6104690 haplotypes had a maximum OR=1.21, indicating that the signal in 20p12.2 can be captured by these two SNPs. The association with MIBC was stronger than with NMIBC for rs6108803 ($OR_{\text{MIBC}}=1.36$, $OR_{\text{NMIBC}}=1.10$, $P_{\text{OR difference}}=0.02$) and rs62185668 ($OR_{\text{MIBC}}=1.39$, $OR_{\text{NMIBC}}=1.13$, $P_{\text{OR difference}}=0.01$) but not for rs6104690 ($OR_{\text{MIBC}}=1.22$, $OR_{\text{NMIBC}}=1.10$, $P_{\text{OR difference}}=0.20$).	Figuroa et al. (2016)
22q13.1 CBX6 (<i>chromobox homolog 6</i>) – APO-BEC3A (<i>apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3A</i>)	Rs1014971[T] (intergenic, 25 kb from CBX6 , 64 kb from APOBEC3A) was associated with UBC in a GWAS (OR=1.14).	Rothman et al. (2010)
	Rs1014971[T] × Smoking interaction (additive) increased significantly UBC risk ($OR_{\text{SNP} \times \text{Smoking}}=2.71$, $P_{\text{additive}}=0.036$).	Garcia-Closas et al. (2013)
	Rs1014971 was not associated with UBC prognosis.	Grotenhuis et al. (2014)
	Rs1014971 UBC risk could not be confirmed in a Chinese study group (1050 cases, 1404 controls, OR=1.14, $P=0.050$).	Wang et al. (2014)

OR: odds ratio, P: P value, HR: hazard ratio, 95 % CI: 95 % confidence interval, adj.: adjusted
 NMIBC: non-muscle invasive bladder cancer, MIBC: muscle-invasive or metastatic bladder cancer,
 PAHs: polycyclic aromatic hydrocarbons

REFERENCES

- Burger M, Catto JWF, Dalbagni G, Grossman HB, Herr H, Karakiewicz P, et al. Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol.* 2013; 63:234-41.
- Carreón T, Hein MJ, Hanley KW, Viet SM, Ruder AM. Bladder cancer incidence among workers exposed to o-toluidine, aniline and nitrobenzene at a rubber chemical manufacturing plant. *Occup Environ Med.* 2014; 71:175-82.
- Delclos GL, Lerner SP. Occupational risk factors. *Scand J Urol Nephrol.* 2008;42:58-63.
- Dudek AM, Grotenhuis AJ, Vermeulen SH, Kienemeny LA, Verhaegh GW. Urinary bladder cancer susceptibility markers. What do we know about functional mechanisms? *Int J Mol Sci.* 2013;14:12346-66.
- Ebbinghaus D, Bánfi G, Selinski S, Blaszkewicz M, Bürger H, Hengstler JG, et al. Polymorphisms of xenobiotic metabolizing enzymes in bladder cancer patients of the Semmelweis University Budapest, Hungary. *J Toxicol Environ Health A.* 2017;80:423-9.
- Figuroa JD, Han SS, Garcia-Closas M, Baris D, Jacobs EJ, Kogevinas M, et al. Genome-wide interaction study of smoking and bladder cancer risk. *Carcinogenesis.* 2014a;35:1737-44.
- Figuroa JD, Ye Y, Siddiq A, Garcia-Closas M, Chatterjee N, Prokunina-Olsson L, et al. Genome-wide association study identifies multiple loci associated with bladder cancer risk. *Hum Mol Genet.* 2014b;23:1387-98.
- Figuroa JD, Middlebrooks CD, Banday AR, Ye Y, Garcia-Closas M, Chatterjee N, et al. Identification of a novel susceptibility locus at 13q34 and refinement of the 20p12.2 region as a multi-signal locus associated with bladder cancer risk in individuals of European ancestry. *Hum Mol Genet.* 2016;25:1203-14.
- Fu YP, Kohaar I, Rothman N, Earl J, Figuroa JD, Ye Y, et al. Common genetic variants in the PSCA gene influence gene expression and bladder cancer risk. *Proc Natl Acad Sci U S A.* 2012;109:4974-9.
- Garcia-Closas M, Malats N, Silverman D, Dosemeci M, Kogevinas M, Hein DW, et al. NAT2 slow acetylation, GSTM1 null genotype, and risk of bladder cancer: results from the Spanish Bladder Cancer Study and meta-analyses. *Lancet.* 2005;366:649-59.
- Garcia-Closas M, Ye Y, Rothman N, Figuroa JD, Malats N, Dinney CP, et al. A genome-wide association study of bladder cancer identifies a new susceptibility locus within SLC14A1, a urea transporter gene on chromosome 18q12.3. *Hum Mol Genet.* 2011;20:4282-9.
- Garcia-Closas M, Rothman N, Figuroa JD, Prokunina-Olsson L, Han SS, Baris D, et al. Common genetic polymorphisms modify the effect of smoking on absolute risk of bladder cancer. *Cancer Res.* 2013;73:2211-20.
- Golka K, Prior V, Blaszkewicz M, Cascorbi I, Schöps W, Kierfeld G, et al. Occupational history and genetic N-acetyltransferase polymorphism in urothelial cancer patients of Leverkusen, Germany. *Scand J Work Environ Health.* 1996;22:332-8.
- Golka K, Reckwitz T, Kempkes M, Cascorbi I, Blaszkewicz M, Reich SE, et al. N-Acetyltransferase 2 (NAT2) and glutathione S-transferase μ (GSTM1) in bladder-cancer patients in a highly industrialized area. *Int J Occup Environ Health.* 1997;3:105-10.
- Golka K, Selinski S, Lehmann ML, Blaszkewicz M, Marchan R, Ickstadt K, et al. Genetic variants in urinary bladder cancer: collective power of the "wimp SNPs". *Arch Toxicol.* 2011;85:539-54.
- Golka K, Prior V, Blaszkewicz M, Bolt HM. The enhanced bladder cancer susceptibility of NAT2 slow acetylators towards aromatic amines: a review considering ethnic differences. *Toxicol Lett.* 2002;128:229-41.
- Golka K, Wiese A, Assennato G, Bolt HM. Occupational exposure and urological cancer. *World J Urol.* 2004;21:382-91.
- Golka K, Hermes M, Selinski S, Blaszkewicz M, Bolt HM, Roth G, et al. Susceptibility to urinary bladder cancer: relevance of rs9642880[T], GSTM1 0/0 and occupational exposure. *Pharmacogenet Genom.* 2009; 19:903-6.
- Golka K, Kopps S, Prager HM, Mende S v, Thiel R, Jungmann O, et al. Bladder cancer in crack testers applying azo dye-based sprays to metal bodies. *J Toxicol Environ Health A.* 2012;75:566-71.
- Grotenhuis AJ, Dudek AM, Verhaegh GW, Witjes JA, Aben KK, van der Marel SL, et al. Prognostic relevance of urinary bladder cancer susceptibility Loci. *PLoS ONE.* 2014;9:e89164.

- Gudbjartsson DF, Helgason H, Gudjonsson SA, Zink F, Oddson A, Gylfason A, et al. Large-scale whole-genome sequencing of the Icelandic population. *Nat Genet.* 2015;47:435-44.
- Höhne S, Gerullis H, Blaszkewicz M, Selinski S, Hengstler JG, Otto T, et al. N-acetyltransferase 1*10 genotype in bladder cancer patients. *J Toxicol Environ Health A.* 2017;80:417-22.
- Kiemeny LA, Thorlacius S, Sulem P, Geller F, Aben KK, Stacey SN, et al. Sequence variant on 8q24 confers susceptibility to urinary bladder cancer. *Nat Genet.* 2008;40:1307-12.
- Kiemeny LA, Sulem P, Besenbacher S, Vermeulen SH, Sigurdsson A, Thorleifsson G, et al. A sequence variant at 4p16.3 confers susceptibility to urinary bladder cancer. *Nat Genet.* 2010;42:415-9.
- Krech E, Selinski S, Blaszkewicz M, Bürger H, Kadhum T, Hengstler JG, et al. Urinary bladder cancer risk factors in an area of former coal, iron, and steel industries in Germany. *J Toxicol Environ Health A.* 2017; 80:430-8.
- Lehmann ML, Selinski S, Blaszkewicz M, Orlich M, Ovsianikov D, Moormann O, et al. Rs710521[A] on chromosome 3q28 close to TP63 is associated with increased urinary bladder cancer risk. *Arch Toxicol.* 2010;84:967-78.
- Lukas C, Selinski S, Prager HM, Blaszkewicz M, Hengstler JG, Golka K. Occupational bladder cancer: Polymorphisms of xenobiotic metabolizing enzymes, exposures, and prognosis. *J Toxicol Environ Health A.* 2017;80:439-52.
- Moore LE, Baris DR, Figueroa JD, Garcia-Closas M, Karagas MR, Schwenn MR, et al. GSTM1 null and NAT2 slow acetylation genotypes, smoking intensity and bladder cancer risk: results from the New England bladder cancer study and NAT2 meta-analysis. *Carcinogenesis.* 2011;32:182-9.
- Nørskov MS, Frikke-Schmidt R, Bojesen SE, Nordestgaard BG, Loft S, Tybjærg-Hansen A. Copy number variation in glutathione-S-transferase T1 and M1 predicts incidence and 5-year survival from prostate and bladder cancer, and incidence of corpus uteri cancer in the general population. *Pharmacogenomics J.* 2011;11: 292-9.
- Ovsianikov D, Selinski S, Lehmann ML, Blaszkewicz M, Moormann O, Haenel MW, et al. Polymorphic enzymes, urinary bladder cancer risk, and structural change in the local industry. *J Toxicol Environ Health A.* 2012;75:557-65.
- Rafnar T, Sulem P, Stacey SN, Geller F, Gudmundsson J, Sigurdsson A, et al. Sequence variants at the TERT-CLPTM1L locus associate with many cancer types. *Nat Genet.* 2009;41:221-7.
- Rafnar T, Vermeulen SH, Sulem P, Thorleifsson G, Aben KK, Witjes JA, et al. European genome-wide association study identifies SLC14A1 as a new urinary bladder cancer susceptibility gene. *Hum Mol Genet.* 2011;20:4268-81.
- Rafnar T, Sulem P, Thorleifsson G, Vermeulen SH, Helgason H, Saemundsdottir J, et al. Genome-wide association study yields variants at 20p12.2 that associate with urinary bladder cancer. *Hum Mol Genet.* 2014;23: 5545-57.
- Rothman N, Garcia-Closas M, Chatterjee N, Malats N, Wu X, Figueroa JD, et al. A multi-stage genome-wide association study of bladder cancer identifies multiple susceptibility loci. *Nat Genet.* 2010;42:978-84.
- Rushton L, Hutchings SJ, Fortunato L, Young C, Evans GS, Brown T, et al. Occupational cancer burden in Great Britain. *Br J Cancer.* 2012;107:S3-7.
- Schwender H, Selinski S, Blaszkewicz M, Marchan R, Ickstadt K, Golka K, et al. Distinct SNP combinations confer susceptibility to urinary bladder cancer in smokers and non-smokers. *PLoS ONE.* 2012;7: e51880. Erratum in *PLoS ONE.* 2015;10:e0137937.
- Selinski S. Genetic variants confer susceptibility to urinary bladder cancer: an updated list of confirmed polymorphisms. *EXCLI J.* 2012;11:743-7.
- Selinski S. Highlight report: functional consequences of urinary bladder cancer risk variants. *EXCLI J.* 2013;12:1017-9.
- Selinski S. Urinary bladder cancer risk variants: recent findings and new challenges of GWAS and confirmatory studies. *Arch Toxicol.* 2014;88:1469-75.
- Selinski S, Blaszkewicz M, Lehmann ML, Ovsianikov D, Moormann O, Guballa C, et al. Genotyping NAT2 with only two SNPs (rs1041983 and rs1801280) outperforms the tagging SNP rs1495741 and is equivalent to the conventional 7-SNP NAT2 genotype. *Pharmacogenet Genomics.* 2011;21:673-8.
- Selinski S, Lehmann ML, Blaszkewicz M, Ovsianikov D, Moormann O, Guballa C, et al. Rs11892031[A] on chromosome 2q37 in an intronic region of the UGT1A locus is associated with urinary bladder cancer risk. *Arch Toxicol.* 2012;86:1369-78.

Selinski S, Blaszkewicz M, Ickstadt K, Hengstler JG, Golka K. Refinement of the prediction of N-acetyltransferase 2 (NAT2) phenotypes with respect to enzyme activity and urinary bladder cancer risk. *Arch Toxicol.* 2013;87:2129-39.

Selinski S, Gerullis H, Otto T, Roth E, Volkert F, Ovsianikov D, et al. Ultra-slow N-acetyltransferase 2 is associated with recurrence-free time in bladder cancer patients. *Eur Urol.* 2017a;71:994-5.

Selinski S, Blaszkewicz M, Ickstadt K, Gerullis H, Otto T, Roth E, et al. Identification and replication of the interplay of four genetic high risk variants for urinary bladder cancer. *Carcinogenesis* 2017b (epub ahead of print). doi: 10.1093/carcin/bgx102.

Tang W, Fu YP, Figueroa JD, Malats N, Garcia-Closas M, Chatterjee N, et al. Mapping of the UGT1A locus identifies an uncommon coding variant that affects mRNA expression and protects from bladder cancer. *Hum Mol Genet.* 2012;21:1918-30.

Wang M, Chu H, Lv Q, Wang L, Yuan L, Fu G, et al. Cumulative effect of genome-wide association study-identified genetic variants for bladder cancer. *Int J Cancer.* 2014;135:2653-60.

Wu X, Ye Y, Kiemeny LA, Sulem P, Rafnar T, Matullo G, et al. Genetic variation in the prostate stem cell antigen gene PSCA confers susceptibility to urinary bladder cancer. *Nat Genet.* 2009;41:991-5. Erratum in *Nat Genet.* 2009;41:1156.