

Letter to the editor:

**LYSOPHOSPHATIDIC ACID IN CARCINOGENESIS AND
TUMOR DEVELOPMENT**

Wiebke Albrecht

IfADo – Leibniz Research Centre for Working Environment and Human Factors at
TU Dortmund, Ardeystr. 67, D-44139 Dortmund, Germany

<http://dx.doi.org/10.17179/excli2018-1638>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License
(<http://creativecommons.org/licenses/by/4.0/>).

Dear Editor,

Recently, Magkrioti and colleagues published a study about the autotaxin-lysophosphatidic acid axis in lung cancer (Magkrioti et al., 2018). Autotaxin (ATX, ENPP2), a secreted glycoprotein that cleaves extracellular lysophosphatidylcholine to generate LPA, is the most relevant factor of extracellular lysophosphatidic acid (LPA) production. LPA is known to activate LPA receptors (LPAR), G-protein coupled receptors that show a widespread distribution and are expressed on many tumor cells (Schleicher et al., 2011; Jonkers and Moolenaar, 2009; Lin et al., 2010; Yu et al., 2016). Magkrioti et al. (2018) demonstrated that a phospholipid phosphatase (LPPP3) that further metabolizes LPA was downregulated in patients with lung cancer, which may lead to increased intratumoral LPA. Moreover, genetic deletion of autotaxin as well as the LPA receptor *Lpar1* attenuated tumor growth in mouse models of lung cancer (Magkrioti et al., 2018).

The role of extracellular LPA in tumor development is well established (Okabe et al., 2011; Leblanc et al., 2018; Stuelten et al., 2018; Bailey et al., 2017; Fukushima et al., 2017). However, recently, evidence has been presented that also intracellular LPA influences the tumor phenotype by enhancing tumor cell migration (Stewart et al., 2012; Hassan, 2017; Lesjak et al., 2014; Marchan et al., 2012). The intracellular glycerophosphocholine phosphodiesterase EDI3 (GDE5, GPD6, or GPCPD1) hydrolyzes the glycerophosphodiester glycerophosphocholine (GPC) to glycerol-3-phosphate (G3P) and choline (Marchan et al., 2017; Stewart et al., 2012). G3P is further metabolized to LPA by glycerol-3-phosphate acyltransferase (GPAM). Recently, it has been shown that knockdown of GPAM decreases, while overexpression increases tumor cell migration (Marchan et al., 2017). These changes in migration correspond to altered intracellular LPA concentrations (Marchan et al., 2017).

It still remains open by which mechanism intracellular LPA influences the phenotype of tumor cells. Does it act on endocytosed LPA receptors? Or are there so far unidentified interaction partners? Tumor development represents a complex process where control factors of proliferation (Schmidt et al., 2008; Cadenas et al., 2014; Hellwig et al., 2016), interactions with the immune system (Schmidt et al., 2012, 2018; Heimes et al., 2017a, b) and redox factors (Cadenas et al., 2010; Jabs et al., 2017) play a key role. Research how phospholipid metabolism contributes to this complex process is still at its infancy.

REFERENCES

- Bailey KA, Klymenko Y, Feist PE, Hummon AB, Stack MS, Schultz ZD. Chemical analysis of morphological changes in lysophosphatidic acid-treated ovarian cancer cells. *Sci Rep.* 2017;7(1):15295.
- Cadenas C, Franckenstein D, Schmidt M, Gehrman M, Hermes M, Geppert B, et al. Role of thioredoxin reductase 1 and thioredoxin interacting protein in prognosis of breast cancer. *Breast Cancer Res.* 2010;12(3):R44.
- Cadenas C, van de Sandt L, Edlund K, Lohr M, Hellwig B, Marchan R, et al. Loss of circadian clock gene expression is associated with tumor progression in breast cancer. *Cell Cycle.* 2014;13:3282-91.
- Fukushima K, Takahashi K, Yamasaki E, O-nishi Y, Fukushima N, Honoki K, Tsujiuchi T. Lysophosphatidic acid signaling via LPA1 and LPA3 regulates cellular functions during tumor progression in pancreatic cancer cells. *Exp Cell Res.* 2017;352:139-45.
- Hassan R. Highlight report: The EDI3-GPAM axis in tumor cell migration. *EXCLI J.* 2017;16:1148-9.
- Heimes AS, Madjar K, Edlund K, Battista MJ, Almstedt K, Elger T, et al. Subtype-specific prognostic impact of different immune signatures in node-negative breast cancer. *Breast Cancer Res Treat.* 2017a;165:293-300.
- Heimes AS, Madjar K, Edlund K, Battista MJ, Almstedt K, Gebhard S, et al. Prognostic significance of interferon regulating factor 4 (IRF4) in node-negative breast cancer. *J Cancer Res Clin Oncol.* 2017b;143:1123-31.
- Hellwig B, Madjar K, Edlund K, Marchan R, Cadenas C, Heimes AS, et al. Epsin family member 3 and ribosome-related genes are associated with late metastasis in estrogen receptor-positive breast cancer and long-term survival in non-small cell lung cancer using a genome-wide identification and validation strategy. *PLoS One.* 2016;11(12):e0167585.
- Jabs V, Edlund K, König H, Grinberg M, Madjar K, Rahnenführer J, et al. Integrative analysis of genome-wide gene copy number changes and gene expression in non-small cell lung cancer. *PLoS One.* 2017;12(11):e0187246.
- Jonkers J, Moolenaar WH. Mammary tumorigenesis through LPA receptor signaling. *Cancer Cell.* 2009;15(6):457-9.
- Leblanc R, Houssin A, Peyruchaud O. Platelets, autotaxin and lysophosphatidic acid signalling: win-win factors for cancer metastasis. *Br J Pharmacol.* 2018;175:3100-10.
- Lesjak MS, Marchan R, Stewart JD, Rempel E, Rahnenführer J, Hengstler JG. EDI3 links choline metabolism to integrin expression, cell adhesion and spreading. *Cell Adh Migr.* 2014;8:499-508.
- Lin S, Lee SJ, Shim H, Chun J, Yun CC. The absence of LPA receptor 2 reduces the tumorigenesis by ApcMin mutation in the in-testine. *Am J Physiol Gastrointest Liver Physiol.* 2010;299:G1128-38.
- Magkrioti C, Oikonomou N, Kaffe E, Mouratis MA, Xylourgidis N, Barbayianni I, et al. The autotaxin-lysophosphatidic acid axis promotes lung carcinogenesis. *Cancer Res.* 2018;78:3634-44.
- Marchan R, Lesjak MS, Stewart JD, Winter R, Seeliger J, Hengstler JG. Choline-releasing glycerophosphodiesterase EDI3 links the tumor metabolome to signaling network activities. *Cell Cycle.* 2012;11:4499-506.
- Marchan R, Büttner B, Lambert J, Edlund K, Glaeser I, Blaszkewicz M, et al. Glycerol-3-phosphate acyltransferase 1 promotes tumor cell migration and poor survival in ovarian carcinoma. *Cancer Res.* 2017;77:4589-601.
- Okabe K, Hayashi M, Yoshida I, Nishimura K, Fukushima N, Tsujiuchi T. Distinct DNA methylation patterns of lysophosphatidic acid receptor genes during rat hepatocarcinogenesis induced by a choline-deficient L-amino acid-defined diet. *Arch Toxicol.* 2011;85:1303-10.
- Schleicher SM, Thotala DK, Linkous AG, Hu R, Leahy KM, Yazlovitskaya EM, et al. Autotaxin and LPA receptors represent potential molecular targets for the radiosensitization of murine glioma through effects on tumor vasculature. *PLoS One.* 2011;6(7):e22182.
- Schmidt M, Böhm D, von Törne C, Steiner E, Puhl A, Pilch H, et al. The humoral immune system has a key prognostic impact in node-negative breast cancer. *Cancer Res.* 2008;68:5405-13.
- Schmidt M, Hellwig B, Hammad S, Othman A, Lohr M, Chen Z, et al. A comprehensive analysis of human gene expression profiles identifies stromal immunoglobulin κ C as a compatible prognostic marker in human solid tumors. *Clin Cancer Res.* 2012;18:2695-703.
- Schmidt M, Weyer-Elberich V, Hengstler JG, Heimes AS, Almstedt K, Gerhold-Ay A, et al. Prognostic impact of CD4-positive T cell subsets in early breast cancer: a study based on the FinHer trial patient population. *Breast Cancer Res.* 2018;20(1):15.

Stewart JD, Marchan R, Lesjak MS, Lambert J, Hergenroeder R, Ellis JK, et al. Choline-releasing glycerophosphodiesterase EDI3 drives tumor cell migration and metastasis. *Proc Natl Acad Sci U S A.* 2012;109:8155-60.

Stuelten CH, Lee RM, Losert W, Parent CA. Lysophosphatidic acid regulates the motility of MCF10CA1a breast cancer cell sheets via two opposing signaling pathways. *Cell Signal.* 2018;45:1-11.

Yu X, Zhang Y, Chen H. LPA receptor 1 mediates LPA-induced ovarian cancer metastasis: an in vitro and in vivo study. *BMC Cancer.* 2016;16:846.