

Original article:

PROTECTIVE EFFECTS OF *SCUTELLARIA BAICALENSIS* GEORGI AGAINST HYDROGEN PEROXIDE-INDUCED DNA DAMAGE AND APOPTOSIS IN HACAT HUMAN SKIN KERATINOCYTES

Jung Jeh Yoon¹, Jin-Woo Jeong¹, Eun Ok Choi¹, Min Ju Kim², Hyun Hwang-Bo¹, Hong Jae Kim¹, Su Hyun Hong¹, Cheol Park³, Dong Hee Lee⁴, Yung Hyun Choi^{1*}

¹ Anti-Aging Research Center and Department of Biochemistry, Dongeui University College of Korean Medicine, 176 Yangjeong-ro, Busanjin-gu, Busan 47227, Republic of Korea

² Lioele Cosmetic Co., LTD., 2068-1 Jungangdae-ro, Geumjeong-gu, Busan 46214, Republic of Korea

³ Department of Molecular Biology, College of Natural Sciences, Dongeui University, 176 Eomgwangno Busanjin-gu, Busan 47340, Republic of Korea

⁴ Genomine Inc., Venture Bldg 306, Pohang TechnoPark, 394 Jigokor Pohang, 37668, Republic of Korea

* Corresponding author: Dr. Y. H. Choi, Department of Biochemistry, Dongeui University College of Korean Medicine, 52-57, Yangjeong-ro, Busanjin-gu, Busan 47227, Republic of Korea, E-mail: choiyh@deu.ac.kr

<http://dx.doi.org/10.17179/excli2016-817>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

Oxidative stress due to excessive accumulation of reactive oxygen species (ROS) is one of the risk factors for the development of several chronic diseases. In this study, we investigated the protective effects of *Scutellaria baicalensis* rhizome ethanol extract (SBRE) against oxidative stress-induced cellular damage and elucidated the underlying mechanisms in the HaCaT human skin keratinocyte cell line. Our results revealed that treatment with SBRE prior to hydrogen peroxide (H₂O₂) exposure significantly increased viability of HaCaT cells. SBRE also effectively attenuated H₂O₂-induced comet tail formation and inhibited the H₂O₂-induced phosphorylation levels of the histone γ H2AX, as well as the number of apoptotic bodies and Annexin V-positive cells. In addition, SBRE exhibited scavenging activity against intracellular ROS generation and restored the mitochondrial membrane potential loss by H₂O₂. Moreover, H₂O₂ enhanced the cleavage of caspase-3 and degradation of poly (ADP-ribose)-polymerase, a typical substrate protein of activated caspase-3, as well as DNA fragmentation; however, these events were almost totally reversed by pretreatment with SBRE. Furthermore, SBRE increased the levels of heme oxygenase-1 (HO-1), which is a potent antioxidant enzyme, associated with the induction of nuclear factor-erythroid 2-related factor 2 (Nrf2). According to our data, SBRE is able to protect HaCaT cells from H₂O₂-induced DNA damage and apoptosis through blocking cellular damage related to oxidative stress through a mechanism that would affect ROS elimination and activating the Nrf2/HO-1 signaling pathway.

Keywords: *Scutellaria baicalensis* rhizome, antioxidant, DNA damage, apoptosis, ROS, Nrf2/HO-1

INTRODUCTION

Accumulating evidence has demonstrated that excessive exposure to ultraviolet (UV) radiation from sunlight is a major risk factor for skin damage due to the overwhelming generation of reactive oxygen species (ROS) (Kammeyer and Luiten, 2015; Karran and Brem, 2016). Although some ROS physiologically act as cellular messengers in redox signaling, uncontrolled release of ROS is further involved in the pathogenesis of a number of human skin disorders by inducing oxidative DNA damage (Birch-Machin et al., 2013; Katiyar, 2016). Consequently, damaged DNA contributes to the initiation of cellular apoptosis in keratinocytes, and this in turn eventually leads to the disruption of the epithelial structure of the skin (Brunelle and Chandel, 2002; Cadet et al., 2015). Therefore, the development of safer and more effective antioxidants for skin protection continues to be an important research goal.

The accumulation of ROS could modulate the expression of redox-sensitive genes. One of the most important regulators in the defense of cells against oxidative stress is nuclear transcription factor erythroid-2-like factor 2 (Nrf2), a basic leucine zipper transcription factor (Satoh et al., 2013; Loboda et al., 2016). This transcription factor is known to activate the expression of antioxidant response element (ARE)-related expression of phase 2-detoxifying genes and anti-oxidants, such as heme oxygenase-1 (HO-1) and NAD(P)H-quinone oxidoreductase-1 (NQO-1) (Satoh et al., 2013; Loboda et al., 2016). Because these antioxidant enzymes are essential for protecting the keratinocytes from oxidative stress (Gęgotek and Skrzydlewska, 2015; Schäfer and Werner, 2015), increasing the antioxidant capacity of skin cells by antioxidant supplements could be a valuable strategy against oxidative stress-induced skin damage.

Scutellaria baicalensis Georgi is a species of flowering plant in the Lamiaceae family, which is widely distributed in Siberia, Mongolia, the Russian Far East, China, and Korea (Li et al., 2011). The root of *S. baicalensis*,

Scutellariae Radix, has been used worldwide in traditional medicine for the treatment of various human diseases (Boyle et al., 2011). Previous studies have indicated that the extracts of *S. baicalensis* and its biogenic active ingredients have various pharmacological properties, including antibacterial (Chung et al., 2014), antioxidant (Lin et al., 2014; Chang et al., 2016), anti-tumor (Choi et al., 2015, 2016a), anti-obesity (Song et al., 2013), anti-asthmatic (Ko et al., 2014; Zhou et al., 2016), anti-hypotensive (Huang et al., 2005; Deng et al., 2012), neuroprotective (Shang et al., 2013; Moon and Park, 2015), and anti-inflammatory activities (Ku and Bae, 2015; Seok et al., 2016). We have recently demonstrated that the protective effects of baicalein, an active component of *S. baicalensis*, against oxidative stress-induced damage through activation of Nrf2 signaling pathway (Choi et al., 2016b). Other researchers have obtained similar results (Zhang et al., 2012; Cui et al., 2015). However, the protective effects and molecular mechanisms involved in oxidative stress-induced skin damages are still unclear. In this study, as a part of our research program to find new antioxidative compounds from the medicinal herbs, we investigated the protective effect of *S. baicalensis* rhizome ethanol extract (SBRE) against hydrogen peroxide (H₂O₂)-induced ROS generation, oxidative damage, and apoptosis in HaCaT immortalized non-tumorigenic human keratinocytes.

MATERIALS AND METHODS

Preparation of SBRE

For the preparation of SBRE, the dried roots of *S. baicalensis* (100 g) were provided by Donggeui Korean Medical Center (Busan, Republic of Korea) and authenticated by Professor S.H. Hong, Department of Biochemistry, Donggeui University College of Korean Medicine. The dried roots (100 g) were cut into small pieces, ground into a fine powder, and then extracted in 1 L of 70 % ethanol by sonication for 3 h. After filtering and concentration of the extracts, the remaining powder was dissolved in dimethyl sulfoxide (DMSO,

Sigma-Aldrich Chemical Co., St. Louis, MO, USA) as a stock solution at a 100 mg/ml concentration and stored at 4 °C. The stock solution was diluted with cell culture medium to the desired concentration prior to use.

Cell culture

HaCaT human skin keratinocyte cell line was obtained from the American Type Culture Collection (Manassas, MD, USA) and cultured in RPMI 1640 medium (WelGENE Inc., Daegu, Republic of Korea) containing 10 % heat-inactivated fetal bovine serum (FBS, WelGENE Inc., Daegu, Republic of Korea), streptomycin (100 µg/mL), and penicillin (100 Units/mL). Cells were maintained at 37 °C in an incubator with a humidified atmosphere of 5 % CO₂.

Cell viability assay

For the cell viability assay, HaCaT cells were seeded in 6-well plate at 2×10^5 cells/ml and cultured for 24 h before being treated with various concentrations of SBRE for 24 h in the presence or absence of H₂O₂ or N-acetylcysteine (NAC, Sigma-Aldrich Chemical Co.). After treatments, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT, Sigma-Aldrich Chemical Co., St. Louis, MO, USA) solution (0.5 mg/ml) was added, followed by 2 h-incubation at 37 °C in the dark, and the medium was removed. The formazan precipitate was dissolved in DMSO. The absorbance of the formazan product was measured at 540 nm using an enzyme-linked immunosorbent assay (ELISA) reader (Molecular Devices, Sunnyvale, CA, USA) (Kim et al., 2016).

Measurement of intracellular ROS

The oxidation-sensitive dye 2',7'-dichlorodihydrofluorescein diacetate (DCF-DA, Molecular Probes, Eugene, OR, USA) was used to determine the formation of intracellular ROS. Briefly, the cells were harvested, washed twice with phosphate-buffered saline (PBS), and then re-suspended in 10 µM DCF-DA for 30 min at 37 °C in the dark. ROS pro-

duction in the cells was monitored immediately by a flow cytometer (Becton Dickinson, San Jose, CA, USA).

Comet assay (Single-cell gel electrophoresis assay)

After each treatment, the cells were washed with PBS and the cell suspension was mixed with 0.5 % low melting agarose (LMA) at 37 °C, and added to the slides pre-coated with 1.0 % normal melting agarose. After solidification of the agarose, slides were covered with another 0.5 % LMA, and then immersed in lysis buffer [2.5 M NaCl, 500 mM Na-ethylenediaminetetraacetic acid (EDTA, Sigma-Aldrich Chemical Co.), 1 M Tris buffer, 1 % sodium lauryl sarcosine, and 1 % Triton X-100) for 1 h at 4 °C. Later, the slides were transferred into an unwinding buffer for another 20 min for DNA unwinding. The slides were then placed in an electrophoresis tank containing 300 mM NaOH and 1 mM Na-EDTA (pH 13.0). An electrical field was then applied (300 mA, 25 V) for 20 min at 25 °C to draw the negatively-charged DNA toward the anode. The slides were washed three times for 5 min at 25 °C in a neutralizing buffer (0.4 M Tris, pH7.5), followed by staining with 20 µg/ml propidium iodide (PI, Sigma-Aldrich Chemical Co.). The slides were examined under a fluorescence microscope.

Protein extraction and Western blot analysis

Total cell lysates were lysed in an extraction buffer [25 mM Tris-Cl (pH 7.5), 250 mM NaCl, 5 mM EDTA, 1 % Nonidet P-40, 0.1 mM sodium orthovanadate, 2 µg/mL leupeptin, and 100 µg/mL phenylmethylsulfonyl fluoride]. The protein concentration in the cell lysate was determined using a detergent-compatible protein assay from Bio-Rad (Hercules, CA, USA). For Western blot analysis, equal amounts of protein (30~50 µg) were separated by 8~10 % sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis and then electro-transferred to polyvinylidene fluoride (PVDF) membranes

(Schleicher & Schuell, Keene, NH, USA). The membranes were blocked with 5 % skim milk for 1 h and then subjected to immunoblot analysis with the appropriate antibodies. Using an enhanced chemiluminescence (ECL) detection system (Amersham Co., Arlington Heights, IL, USA), immunoreactive bands were detected and exposed to X-ray film. Various primary and secondary antibodies for Western blot analysis were obtained from Cell Signaling Technology, Inc. (Boston, MA, USA) and Santa Cruz Biotechnology (Santa Cruz, CA, USA).

Measurement of Mitochondrial Membrane Potential (MMP, $\Delta\psi_m$)

MMP values were determined using the dual-emission potential-sensitive probe 6,6'-tetrachloro-1,1',3,3'-tetraethylimidacarbocyanine iodide (JC-1, Sigma-Aldrich Chemical Co.). Briefly, the cells were collected and incubated with 10 μ M of JC-1 for 20 min at 37 °C in the dark. After JC-1 was removed, the cells were washed with PBS to remove unbound dye, and the amount of JC-1 retained by 10,000 cells per sample was measured using a flow cytometer (Chun et al., 2016).

DNA fragmentation assay

For the extraction of fragmented DNAs, the cells were lysed in a buffer [10 mM Tris-HCl (pH 7.4), 150 mM NaCl, 5 mM EDTA, and 0.5 % Triton X-100] for 1 h at room temperature. After centrifugation at 14,000 rpm for 30 min, the supernatant was collected and incubated with proteinase K (Invitrogen, Carlsbad, CA, USA) for 3 h at 50 °C. The fragmented DNA in the supernatant was separated through 1 % agarose gel. DNA fragmentation pattern was visualized by ultraviolet light source after ethidium bromide (EtBr, Sigma-Aldrich Chemical Co.) staining.

Nuclear staining with DAPI

The cells were harvested, washed with PBS, and fixed with 3.7 % paraformaldehyde (Sigma-Aldrich Chemical Co.) for 30 min at room temperature. After washing with PBS twice, cells were stained with 2.5 μ g/ml 4,6-

dianmidino-2-phenylindole (DAPI, Sigma-Aldrich Chemical Co.) solution for 20 min at room temperature. The stained cells were washed three times with PBS and analyzed using a fluorescence microscope (Carl Zeiss, Deisenhofen, Germany).

Determination of apoptotic cells by flow cytometry

A fluorescein-conjugated annexin V (annexin V-FITC) staining assay kit (BD Biosciences, San Jose, CA, USA) was used to quantitatively assess the level of induced cell apoptosis. Briefly, the cells were washed with PBS and stained with 5 μ l of Annexin V-FITC and 5 μ l of propidium iodide (PI) in each sample to quantify the cell number at different stages of cell death. After 15 min incubation at room temperature in the dark, the degree of apoptosis was quantified as a percentage of the Annexin V-positive and PI-negative (Annexin V⁺/PI⁻ cells) cells using a flow cytometer.

Statistical analysis

All experiments were conducted in triplicate, and a one-way ANOVA test (SPSS 11.5 statistical software) was used to compare the mean values of each treatment. Significant differences were determined using the Duncan's multiple range test. *P < 0.05 was considered to be significant.

RESULTS

SBRE protects the HaCaT keratinocytes from H₂O₂-induced cytotoxicity

Prior to investigating the protective properties of SBRE against H₂O₂ treatment, the cytotoxic effects of SBRE were first examined on HaCaT cells. The MTT assay indicated that the treatments did not result in any cytotoxic effect up to the concentration of 600 μ g/ml, whereas cell viability significantly decreased at concentrations higher than 800 μ g/ml (Figure 1A). Thus, 600 μ g/ml of SBRE was selected as the optimum concentration for the subsequent examination of the protective effect of SBRE on H₂O₂-induced

cytotoxicity. Further MTT assays revealed that treatment with 1 mM H₂O₂ significantly reduced cell viability; however, H₂O₂-induced reduction of cell viability was effectively protected by pretreatment with SBRE in a concentration-dependent manner. A pretreatment with NAC, a potent antioxidant, also effectively abrogated H₂O₂-induced cytotoxicity (Figure 1B).




Figure 1: Effects of SBRE on the H₂O₂-induced growth inhibition in HaCaT keratinocyte cells. The cells were treated with the indicated concentrations of SBRE for 24 h (A) or pre-treated with or without SBRE (200, 400 and 600 µg/ml) and NAC (10 mM) for 1 h, then cultured with H₂O₂ (1 mM) for 24 h (B). The cell viability was measured by MTT assay. The results are the mean ± SD values obtained in three independent experiments (*P < 0.05 significant as compared to the H₂O₂-treated group; #P < 0.05 vs. H₂O₂-treated cells).

SBRE suppresses H₂O₂-induced ROS generation in HaCaT keratinocytes

To study the mechanisms underlying the protective effect of SBRE, the radical-scavenging effect of SBRE was determined through the use of an ROS-sensitive fluorescent dye, DCF-DA. The results of the flow cytometry analysis demonstrated that HaCaT

cells exposed to H₂O₂ for 30 min showed a significant increase in the generation of ROS, whereas this induction was substantially inhibited by SBRE or NAC pretreatment (Figure 2). The results indicated that the protective effects of SBRE against H₂O₂ in HaCaT cells were a direct consequence of the inhibition of excessive ROS production.

SBRE inhibits H₂O₂-induced DNA damage in HaCaT keratinocytes

We next assessed the protective effects of SBRE against H₂O₂-induced DNA damages using the comet assay, which measures single as well as double-strand breaks (Azqueta et al., 2014). Figure 3 demonstrates that H₂O₂ treatment increased both the amounts of DNA in the tail (tail moment) and distance of DNA migration (tail length); however, SBRE pretreatment significantly reduced both percentages elevated by H₂O₂. Additionally, immunoblotting results revealed that exposure of cells to H₂O₂ increased the phosphorylation of histone γH2AX on serine 139, a marker of DNA double strand breaks (Rogakou et al., 1998); however, this adverse effect was effectively inhibited by SBRE pretreatment (Figure 4). Our findings suggested that inhibition of excessive ROS by SBRE may contribute to the attenuation of H₂O₂-induced DNA damage.

SBRE alleviates H₂O₂-induced mitochondrial dysfunction in HaCaT keratinocytes

A change in MMP ($\Delta\psi_m$) was examined because mitochondria are the major sites of oxidative phosphorylation and production of ROS, and they are involved in the initiation of apoptosis through membrane permeabilization (Cha et al., 2015). As shown in Figure 5, H₂O₂-treated cells showed a loss of MMP, as substantiated by an increase in fluorescence (FL-1) with the JC-1 dye in flow cytometry analysis, indicative of mitochondrial depolarization, as compared with control cells and SBRE-treated cells. However, pretreatment with SBRE significantly blocked the loss of MMP in H₂O₂-treated cells as did NAC.




Figure 2: Effects of SBRE on H₂O₂-induced ROS generation in HaCaT keratinocyte cells. Cells were pretreated with 600 µg/ml SBRE or 10 mM NAC for 1 h and then stimulated with or without 1 mM H₂O₂ for 30 min. The cells were incubated at 37 °C in the dark for 20 min with culture medium containing 10 µM DCF-DA to monitor ROS production. **(A)** ROS generation was measured by flow cytometry. **(B)** Each point represents the mean ± SD of three independent experiments (*P < 0.05 vs. untreated control; #P < 0.05 vs. H₂O₂-treated cells).




Figure 3: Protection of H₂O₂-induced DNA damage by SBRE in HaCaT keratinocyte cells. Cells were pretreated with 600 µg/ml SBRE for 1 h and then incubated with or without 1 mM H₂O₂ for 24 h. **(A)** To detect cellular DNA damage, the comet assay was performed, and representative pictures of the comets were taken using a fluorescence microscope (original magnification, 400×). **(B and C)** The average of the Tail Moment (TM) and Tail Length of at least 100 cells per experimental point are shown. (*P < 0.05 significant as compared to untreated control group; #P < 0.05 significant as compared to H₂O₂-treated group).




Figure 4: Attenuation of H₂O₂-induced phosphorylation of γH2AX by SBRE in HaCaT keratinocyte cells. Cells were pretreated with 600 µg/ml SBRE for 1 h and then stimulated with or without 1 mM µM H₂O₂ for 24 h. **(A)** The cells were lysed and then equal amounts of cell lysates were separated on SDS-polyacrylamide gels and transferred to PVDF membranes. The membranes were probed with specific antibodies against γH2AX and p-γH2AX, and the proteins were visualized using an ECL detection system. Actin was used as an internal control. **(B)** The relative expression of p-γH2AX represents the average densitometric analyses as compared with γH2AX. Each point represents the mean ± SD of three independent experiments (*P < 0.05 vs. untreated control; #P < 0.05 vs. H₂O₂-treated cells).




Figure 5: Inhibition of H₂O₂-induced mitochondrial dysfunction by SBRE in HaCaT keratinocyte cells. Cells were pretreated with 600 µg/ml SBRE or 10 mM NAC for 1 h and then stimulated with or without 1 mM H₂O₂ for 24 h. **(A)** The cells were collected and incubated with 10 µM JC-1 for 20 min at 37 °C in the dark. The cells were then washed once with PBS, and the values of MMP were evaluated using a flow cytometer. **(B)** Each point represents the mean ± SD of three independent experiments (*P < 0.05 vs. untreated control; #P < 0.05 vs. H₂O₂-treated cells).

SBRE prevents H₂O₂-induced apoptosis in HaCaT keratinocytes

To elucidate the cytoprotective effect of SBRE against the H₂O₂-induced reduction of HaCaT cell viability, it was investigated the effects of SBRE on H₂O₂-mediated apoptosis. The results of nuclear morphological changes and DNA fragmentation using DAPI staining and agarose gel electrophoresis, respectively, showed that H₂O₂ treatment significantly increased the number of condensed or blebbing nuclei and formation of DNA laddering, whereas, when these cells were pretreated with SBRE, these phenomena were markedly reduced (Figure 6A-C). The results of flow cytometry analysis consistently indicated an increased number of annexin-positive cells, i.e., apoptotic cells, in the H₂O₂-treated culture compared to the untreated control group. However, the pretreatment of cells with SBRE prior to H₂O₂ exposure effectively protected the cells against apoptosis (Figure 7).

whereas, when these cells were pretreated with SBRE, these phenomena were markedly reduced (Figure 6A-C). The results of flow cytometry analysis consistently indicated an increased number of annexin-positive cells, i.e., apoptotic cells, in the H₂O₂-treated culture compared to the untreated control group. However, the pretreatment of cells with SBRE prior to H₂O₂ exposure effectively protected the cells against apoptosis (Figure 7).




Figure 6: Prevention of H₂O₂-induced chromatin condensation, DNA fragmentation, cleavage of caspase-3 and degradation of PARP by SBRE in HaCaT keratinocyte cells. Cells were pretreated with 600 µg/ml for 1 h and then stimulated with or without 1 mM µM H₂O₂ for 24 h. **(A and B)** The cells were fixed and stained with DAPI solution. After 15 min incubation at room temperature, stained nuclei were observed using a fluorescence microscope (original magnification, 400x). Each point represents the mean ± SD of three independent experiments (*P < 0.05 vs. untreated control; #P < 0.05 vs. H₂O₂-treated cells). **(C)** DNA fragmentation was analyzed by extracting the fragmented DNAs and separating them by electrophoresis in a 1.0 % agarose gel containing EtBr. **(D)** The cells were lysed and then equal amounts of cell lysates were separated on SDS-polyacrylamide gels and transferred to nitrocellulose membranes. The membranes were probed with specific antibodies against caspase-3 and PARP, and the proteins were visualized using an ECL detection system. Actin was used as an internal control.




Figure 7: Attenuation of H₂O₂-induced apoptosis by SBRE in HaCaT keratinocyte cells. Cells were pre-treated with 600 µg/ml SBRE or 10 mM NAC for 1 h and then stimulated with or without 1 mM H₂O₂ for 24 h. **(A)** The cells were stained with FITC-conjugated Annexin-V and PI for flow cytometry analysis. The percentages of apoptotic cells were determined by counting the percentage of Annexin V-positive cells. **(B)** Each point represents the mean ± SD of three independent experiments (*P < 0.05 vs. untreated control; #P < 0.05 vs. H₂O₂-treated cells).

SBRE attenuates H₂O₂-induced activation of caspase-3 and cleavage of PARP in HaCaT keratinocytes

The mitochondria-mediated apoptosis pathway is initiated by the loss of MMP and subsequent release of pro-apoptotic proteins, leading to the activation of caspase-9 and -3 (MacKenzie and Clark, 2008; Hensley et al., 2013). Accordingly, the active forms of caspase-3 and degradation of poly(ADP-ribose) polymerase (PARP), a well-known substrate of caspase-3 (Lazebnik et al., 1994), were assessed by Western blot analysis. Our results indicated that there was a marked increase in the levels of activated caspase-3 (17 and 19 kDa) and cleaved PARP (89 kDa) expression in H₂O₂-treated cells compared with

the control (Figure 6D). However, H₂O₂-induced caspase-3 activation and PARP degradation were absolutely abrogated by SBRE pretreatment, suggesting that SBRE protects HaCaT cells from apoptosis by inhibiting the mitochondrial caspase-dependent pathway.

SBRE enhances the expression Nrf2 and HO-1 in HaCaT keratinocytes

A transcription factor Nrf2 regulates the expression of ARE-driven antioxidant and cytoprotective genes including HO-1 and NQO-1 to control the cellular defense against oxidative stress (Satoh et al., 2013; Murakami and Motohashi, 2015). Therefore, we further investigated whether SBRE activated Nrf2, and found that SBRE treatment gradually in-

creased Nrf2 expression as well as its phosphorylation in a time-dependent manner, and concomitantly decreased Kelch-like ECH-associated protein 1 (Keap1) levels, a negative regulator of Nrf2 (Figure 8). Additionally, SBRE also enhanced HO-1 expression, but not NQO-1 expression, indicating that the cytoprotective effects of SBRE against oxidative stress in HaCaT cells are probably associated with activation of the Nrf2/HO-1 signaling pathway.




Figure 8: Induction of Nrf2 and HO-1 expression by SBRE in HaCaT keratinocyte cells. Cells were incubated with 600 µg/ml SBRE for the indicated periods. Total cellular proteins were separated on SDS-polyacrylamide gels and then transferred onto PVDF membranes. The membranes were probed with the indicated antibodies. Proteins were visualized using an ECL detection system. Actin was used as an internal control of total cellular and nuclear proteins.

DISCUSSION

In the present study, we evaluated the antioxidant capacity of SBRE, an ethanol extract of the root of *S. baicalensis*, to prevent oxidative stress-induced cellular damage in HaCaT human skin keratinocytes. Our data from the MTT assay and flow cytometry demonstrated that SBRE significantly rescued cell viability and reduced apoptosis from H₂O₂-treated HaCaT cells. The protection against oxidative stress was mediated by the

inhibition of DNA damage, mitochondrial dysfunction, ROS generation and caspase-3 activation. This protective effect of SBRE was also associated with the activation of the Nrf2/HO-1 signaling pathway.

Oxidative stress, represented as an increase in ROS, is an abnormal phenomenon characterized by a production of free radicals that exceeds the antioxidant capacity. The extremely elevated levels of ROS can destroy cytoprotective defense mechanisms by the neutralizing of antioxidants, leading to the acceleration of skin aging and development of various skin diseases (Kammeyer and Luiten, 2015; Karran and Brem, 2016). Therefore, prevention of ROS accumulation with antioxidant agents is often regarded as a possible strategy for the management of such oxidative damage in the skin (Oresajo et al., 2012; Tundis et al., 2015). The generation of excess ROS can irreversibly cause a range of base modification and DNA strand breaks, resulting in DNA damage (Birch-Machin et al., 2013; Katiyar, 2016). Elevated levels of ROS are associated with the first events in the apoptosis by inducing mitochondrial dysfunction, resulting in the decrease of MMP, and thus activate caspase-9 and -3 through release of mitochondrial apoptotic factors into cytoplasm, which are known to be strong stimulators of the intrinsic apoptosis pathway involving the mitochondrial cascade (Brunelle and Chandel, 2002; Cadet et al., 2015). The activation of caspase-3 participates in the cleavage or degradation of various important proteins including PARP, which are involved in apoptosis, and eventually in the activation of other caspases. Thus, activation of caspase-3 is associated with the cleavage of PARP and has a primary role in triggering the cascade of events leading to the apoptosis pathway (MacKenzie and Clark, 2008; Hensley et al., 2013). In the present study, we demonstrated that SBRE significantly protected the HaCaT skin keratinocyte cells against H₂O₂-induced growth inhibition and DNA damage, suggesting that SBRE may potentially enhance DNA repair. Additionally, along with elevated ROS and mitochondrial dysfunction, we found the

loss of MMP, activation of caspase-3 and degradation of PARP in HaCaT cells by H₂O₂ treatment. However, the results of JC-1 staining and immunoblotting showed SBRE could effectively restore the H₂O₂-induced loss of MMP to the basal level and prevent activation of caspase-3 as well as cleavage of PARP by H₂O₂ in HaCaT cells. These results pointed out that the ability of SBRE to attenuate oxidative stress partly depends on inhibiting mitochondrial-related apoptosis. We also found that elevated ROS accumulation in H₂O₂-exposed cells was substantially inhibited by SBRE pretreatment, indicating the free radical scavenging activity and protective properties of SBRE.

Accumulating evidence strongly suggests that the Nrf2-mediated signaling pathway is essential in protecting human skin fibroblasts against oxidative stress (Satoh et al., 2013; Loboda et al., 2016). Under basal conditions, Nrf2-dependent transcription is suppressed by Keap1, a negative regulator of Nrf2, which facilitates the degradation of Nrf2 through ubiquitin-mediated proteasomal degradation (O'Connell et al., 2015; Jaramillo and Zhang, 2013). Upon modification of specific thiols by insult, Keap1 triggers the dissociation of Nrf2 from the Nrf2-Keap1 complex in the cytoplasm and allows Nrf2 to translocate to the nucleus, where it subsequently activates AREs present in the promoter regions of an array of genes (Satoh et al., 2013; Murakami and Motohashi, 2015). Therefore, the status of Nrf2 and its inhibitory protein Keap-1 determines the Nrf2-mediated ARE activity (Suzuki and Yamamoto, 2015; Loboda et al., 2016). Moreover, several lines of studies revealed that phosphorylation of Nrf2 (Try568) leads to the nuclear export of Nrf2 (Jain and Jaiswal, 2006; Kaspar and Jaiswal, 2011). Therefore, we investigated whether Nrf2 pathway contributes to the protective effects of SBRE against H₂O₂-induced oxidative stress in HaCaT cells. Our data strongly supports the ability of SBRE to stimulate the Nrf2 pathway, as SBRE treatment time-dependently increased Nrf2 accumulation and

phosphorylation, and decreased Keap1 expression. Following the treatment of HaCaT cells with SBRE, we also observed a marked increase in HO-1 expression, but NQO-1 was unaffected by SBRE treatment. HO-1 is a Nrf2 downstream target and powerful indirect antioxidant enzyme, which is a rate-limiting enzyme in heme catabolism (Srisook et al., 2005; Chau, 2015). This enzyme converts heme to beneficial byproducts such as carbon monoxide and bilirubin that can directly scavenge free radicals and repair DNA damage caused by oxidative stress (Jaramillo and Zhang, 2013; Chau, 2015). Although the detailed molecular mechanism of SBRE-mediated Nrf2 activation was not yet fully established, our observations support the proposal that the cytoprotective effect of SBRE against oxidative stress in HaCaT cells is probably mediated through activating the Nrf2/HO-1 signaling pathway. Additionally, all the effects of SBRE were observed at nontoxic concentrations. Therefore SBRE could be considered as important natural active principles to include among skin photoprotective agents with promising applications in dermatological clinical research.

In conclusion, SBRE is a potent antioxidant, able to prevent oxidative DNA damage and to reduce ROS production and activation of the mitochondria-mediated apoptotic pathway in H₂O₂-treated HaCaT human skin keratinocyte cells. This process is also associated with the involvement of Nrf2 activation and the up-regulation of the expression of its downstream antioxidant gene HO-1 in order to protect cells from oxidative stress. Therefore, SBRE may be of therapeutic value in the prevention and treatment of various human skin diseases associated with oxidative stress. However, further studies especially those using human systems are needed to determine cellular uptake, distribution and long-term effects of SBRE on the skin for optimal protection.

Conflicts of interest

The authors declare that there is no conflict of interest.

Acknowledgement

This research was financially supported by the Ministry of Trade, Industry, and Energy (MOTIE), Korea, under the “Regional Specialized Industry Development Program” supervised by the Korea Institute for Advancement of Technology (KIAT, R0003671 and R0005567).

REFERENCES

- Azqueta A, Slyskova J, Langie SA, O'Neill Gaivão I, Collins A. Comet assay to measure DNA repair: approach and applications. *Front Genet.* 2014;5:288.
- Birch-Machin MA, Russell EV, Latimer JA. Mitochondrial DNA damage as a biomarker for ultraviolet radiation exposure and oxidative stress. *Br J Dermatol.* 2013;169(Suppl 2):9-14.
- Boyle SP, Doolan PJ, Andrews CE, Reid RG. Evaluation of quality control strategies in *Scutellaria* herbal medicines. *J Pharm Biomed Anal.* 2011;54:951-7.
- Brunelle JK, Chandel NS. Oxygen deprivation induced cell death: an update. *Apoptosis.* 2002;7:475-82.
- Cadet J, Douki T, Ravanat JL. Oxidatively generated damage to cellular DNA by UVB and UVA radiation. *Photochem Photobiol.* 2015;91:140-55.
- Cha MY, Kim DK, Mook-Jung I. The role of mitochondrial DNA mutation on neurodegenerative diseases. *Exp Mol Med.* 2015;47:e150.
- Chang WS, Lin EY, Hsu SW, Hu PS, Chuang CL, Liao CH, et al. Baicalin scavenged reactive oxygen species and protected human keratinocytes against UVB-induced cytotoxicity. *In Vivo.* 2016;30:605-10.
- Chau LY. Heme oxygenase-1: emerging target of cancer therapy. *J Biomed Sci.* 2015;22:22.
- Choi BB, Choi JH, Park SR, Kim JY, Hong JW, Kim GC. *Scutellariae radix* induces apoptosis in chemo-resistant SCC-25 human tongue squamous carcinoma cells. *Am J Chin Med.* 2015;43:167-81.
- Choi EO, Park C, Hwang HJ, Hong SH, Kim GY, Cho EJ, et al. Baicalein induces apoptosis via ROS-dependent activation of caspases in human bladder cancer 5637 cells. *Int J Oncol.* 2016a;49:1009-18.
- Choi EO, Jeong JW, Park C, Hong SH, Kim GY, Hwang HJ, et al. Baicalein protects C6 glial cells against hydrogen peroxide-induced oxidative stress and apoptosis through regulation of the Nrf2 signaling pathway. *Int J Mol Med.* 2016b;37:798-806.
- Chun SK, Go K, Yang MJ, Zendejas I, Behrns KE, Kim JS. Autophagy in ischemic livers: A critical role of sirtuin 1/mitofusin 2 axis in autophagy induction. *Toxicol Res.* 2016;32:35-46.
- Chung ST, Chien PY, Huang WH, Yao CW, Lee AR. Synthesis and anti-influenza activities of novel baicalein analogs. *Chem Pharm Bull (Tokyo).* 2014;62:415-21.
- Cui G, Luk SC, Li RA, Chan KK, Lei SW, Wang L, et al. Cytoprotection of baicalein against oxidative stress-induced cardiomyocytes injury through the Nrf2/Keap1 pathway. *J Cardiovasc Pharmacol.* 2015;65:39-46.
- Deng YF, Aluko RE, Jin Q, Zhang Y, Yuan LJ. Inhibitory activities of baicalin against renin and angiotensin-converting enzyme. *Pharm Biol.* 2012;50:401-6.
- Gęgotek A, Skrzydlewska E. The role of transcription factor Nrf2 in skin cells metabolism. *Arch Dermatol Res.* 2015;307:385-96.
- Hensley P, Mishra M, Kyprianou N. Targeting caspases in cancer therapeutics. *Biol Chem.* 2013;394:831-43.
- Huang Y, Tsang SY, Yao X, Chen ZY. Biological properties of baicalein in cardiovascular system. *Curr Drug Targets Cardiovasc Haematol Disord.* 2005;5:177-84.
- Jain AK, Jaiswal AK. Phosphorylation of tyrosine 568 controls nuclear export of Nrf2. *J Biol Chem.* 2006;281:12132-42.
- Jaramillo MC, Zhang DD. The emerging role of the Nrf2-Keap1 signaling pathway in cancer. *Genes Dev.* 2013;27:2179-91.
- Kammeyer A, Luiten RM. Oxidation events and skin aging. *Ageing Res Rev.* 2015;21:16-29.
- Karran P, Brem R. Protein oxidation, UVA and human DNA repair. *DNA Repair (Amst).* 2016;44:178-85.
- Kaspar JW, Jaiswal AK. Tyrosine phosphorylation controls nuclear export of Fyn, allowing Nrf2 activation of cytoprotective gene expression. *FASEB J.* 2011;25:1076-87.
- Katiyar SK. Dietary proanthocyanidins inhibit UV radiation-induced skin tumor development through functional activation of the immune system. *Mol Nutr Food Res.* 2016;60:1374-82.
- Kim SH, Kang SH, Kang BS. Therapeutic effects of dihydroartemisinin and transferrin against glioblastoma. *Nutr Res Pract.* 2016;10:393-7.

- Ko WH, Huang Y. Effect of *Scutellariae radix* (Huangqin) on preventing rhinovirus-provoked asthmatic inflammation in cultured human bronchial epithelia. *Hong Kong Med J*. 2014;20(Suppl 4):34-6.
- Ku SK, Bae JS. Baicalin, baicalein and wogonin inhibits high glucose-induced vascular inflammation *in vitro* and *in vivo*. *BMB Rep*. 2015;48:519-24.
- Lazebnik YA, Kaufmann SH, Desnoyers S, Poirier GG, Earnshaw WC. Cleavage of poly(ADP-ribose) polymerase by a proteinase with properties like ICE. *Nature*. 1994;371:346-7.
- Li C, Lin G, Zuo Z. Pharmacological effects and pharmacokinetics properties of *Radix Scutellariae* and its bioactive flavones. *Biopharm Drug Dispos*. 2011;32:427-45.
- Lin M, Li L, Zhang Y, Zheng L, Xu M, Rong R, et al. Baicalin ameliorates H₂O₂ induced cytotoxicity in HK-2 cells through the inhibition of ER stress and the activation of Nrf2 signaling. *Int J Mol Sci*. 2014;15:12507-22.
- Loboda A, Damulewicz M, Pyza E, Jozkowicz A, Dulak J. Role of Nrf2/HO-1 system in development, oxidative stress response and diseases: an evolutionarily conserved mechanism. *Cell Mol Life Sci*. 2016;73:3221-47.
- MacKenzie SH, Clark AC. Targeting cell death in tumors by activating caspases. *Curr Cancer Drug Targets*. 2008;8:98-109.
- Moon JH, Park SY, Baicalein prevents human prion protein-induced neuronal cell death by regulating JNK activation. *Int J Mol Med*. 2015;35:439-45.
- Murakami S, Motohashi H. Roles of Nrf2 in cell proliferation and differentiation. *Free Radic Biol Med*. 2015;88:168-78.
- O'Connell MA, Hayes JD. The Keap1/Nrf2 pathway in health and disease: from the bench to the clinic. *Biochem Soc Trans*. 2015;43:687-9.
- Oresajo C, Pillai S, Manco M, Yatskayer M, McDaniel D. Antioxidants and the skin: understanding formulation and efficacy. *Dermatol Ther*. 2012;25: 252-9.
- Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem*. 1998;273:5858-68.
- Satoh T, McKercher SR, Lipton SA. Nrf2/ARE-mediated antioxidant actions of pro-electrophilic drugs. *Free Radic Biol Med*. 2013;65:645-57.
- Schäfer M, Werner S. Nrf2-A regulator of keratinocyte redox signaling. *Free Radic Biol Med*. 2015;88:243-52.
- Seok JK, Kwak JY, Choi GW, An SM, Kwak JH, Seo HH, et al. *Scutellaria radix* extract as a natural UV protectant for human skin. *Phytother Res*. 2016;30: 374-9.
- Shang Y, Zhang H, Cheng J, Miao H, Liu Y, Cao K, et al. Flavonoids from *Scutellaria baicalensis* Georgi are effective to treat cerebral ischemia/reperfusion. *Neural Regen Res*. 2013;8:514-22.
- Song KH, Lee SH, Kim BY, Park AY, Kim JY. Extracts of *Scutellaria baicalensis* reduced body weight and blood triglyceride in db/db mice. *Phytother Res*. 2013;27:244-50.
- Srisook K, Kim C, Cha YN. Molecular mechanisms involved in enhancing HO-1 expression: de-repression by heme and activation by Nrf2, the "one-two" punch. *Antioxid Redox Signal*. 2005;7:1674-87.
- Suzuki T, Yamamoto M. Molecular basis of the Keap1-Nrf2 system. *Free Radic Biol Med*. 2015;88:93-100.
- Tundis R, Loizzo MR, Bonesi M, Menichini F. Potential role of natural compounds against skin aging. *Curr Med Chem*. 2015;22:1515-38.
- Zhang Z, Cui W, Li G, Yuan S, Xu D, Hoi MP, et al. Baicalein protects against 6-OHDA-induced neurotoxicity through activation of Keap1/Nrf2/HO-1 and involving PKC α and PI3K/AKT signaling pathways. *J Agric Food Chem*. 2012;60:8171-82.
- Zhou DG, Diao BZ, Zhou W, Feng JL. Oroxylin A inhibits allergic airway inflammation in ovalbumin (OVA)-induced asthma murine model. *Inflammation*. 2016;39:867-72.