

Original article:

**ANTI-INFLAMMATORY EFFECT OF SUPERCRITICAL EXTRACT
AND ITS CONSTITUENTS FROM *ISHIGE OKAMURAE***

Eun-Yi Ko^{1,2,a}, Weon-Jong Yoon^{3,a}, Hae-Won Lee⁴, Soo-Jin Heo⁵, Young-Hwan Ko⁶,
I.P. Shanura Fernando², Kichul Cho¹, Chi-Heon Lee^{1,2}, Sung-Pyo Hur⁵, Su-Hyeon Cho^{1,2},
Ginnae Ahn⁷, Daekyung Kim^{1,*}, Kil-Nam Kim^{1,8,*}

¹ Jeju Center, Korea Basic Science Institute (KBSI), Jeju 690-140, Republic of Korea

² Department of Marine Life Science, Jeju National University, Jeju 690-756, Republic of Korea

³ Jeju Biodiversity Research Institute, Jeju Technopark, Jeju, 699-943, Republic of Korea

⁴ World Institute of Kimchi, Gwangju 503-360, Republic of Korea

⁵ Jeju International Marine Science Center for Research & Education, Korea Institute of Ocean Science & Technology (KIOST), Jeju, 63349, Republic of Korea

⁶ Department of Food Bioengineering, Jeju National University, Jeju 690-756, Republic of Korea

⁷ Department of Marine Bio-Food Sciences, Chonnam National University, Yeosu 550-74, Republic Korea

⁸ Department of Marine Biotechnology, University of Science and Technology, Daejeon 305-350, Republic of Korea

* Corresponding authors:

Tel: +82-64-800-4930, E-mail: dkim@kbsi.re.kr (D. Kim)

Tel: +82-64-800-4933, E-mail: knkim@kbsi.re.kr (K.N. Kim)

^a These authors contributed equally to this work

<http://dx.doi.org/10.17179/excli2016-337>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

The anti-inflammatory properties of the supercritical fluid extract of *Ishige okamurae* (SFEIO) on lipopolysaccharide (LPS)-stimulated murine RAW 264.7 macrophages. The lipid profile of the SFEIO, revealed the presence of palmitic acid (220.2 mg/g), linoleic acid (168.0 mg/g), and oleic acid (123.0 mg/g). SFEIO was found to exert its anti-inflammatory effects through inhibiting nitric oxide (NO), prostaglandin E₂ (PGE₂), inducible nitric oxide synthase (iNOS), cyclooxygenase-2 (COX-2), tumor necrosis factor (TNF)- α , interleukin (IL)-1 β , and IL-6 production in LPS-stimulated RAW 264.7 cells, without inducing cytotoxicity. SFEIO did not effect on the LPS-induced p38 kinase phosphorylation, whereas it attenuated the extracellular-related signaling kinase (ERK) and c-Jun N-terminal kinase (JNK) phosphorylation. Furthermore, SFEIO inhibited the LPS-induced I κ B- α degradation and p50 NF- κ B activation. These results suggest that SFEIO exerts its anti-inflammatory effects in LPS-activated RAW 264.7 cells by down-regulating the activation of ERK, JNK, and NF- κ B.

Keywords: supercritical fluid extraction, linoleic acid, *Ishige okamurae*, anti-inflammatory, NF- κ B

INTRODUCTION

Inflammatory diseases have become one of the major health issues in many parts of the world having a considerable influence on the physiological well-being and health care costs. Inflammation can be defined as a non-specific protective response of the body to defend the host from pathogens, toxins, and local injuries. Inflammatory reactions are typically characterized by the redness, swelling, heat, and pain (Medzhitov, 2008). Pathogen- and host-derived molecules (lipopolysaccharide (LPS) and interferon- γ (IFN- γ)) can stimulate macrophages, thereby upregulate inflammatory mediators. These includes nitric oxide (NO), prostaglandin E₂ (PGE₂), inducible nitric oxide synthase (iNOS), and cyclooxygenase-2 (COX-2) (Korhonen et al., 2005). These are considered as some of the major inflammatory mediators that regulate essential steps in inflammatory signaling pathways. Therefore, inhibiting the production of these inflammatory mediators is an important target for the anti-inflammatory drugs to treat inflammatory diseases.

Among a countless number of extraction methods in natural product chemistry, carbon dioxide (CO₂) has gained much attention as an extraction solvent for natural products. It is a non-toxic, readily available and non-explosive solvent that could easy to remove from the extract. The extraction can be achieved by the supercritical fluid extraction (SFE) methodology which utilizes low temperatures, resulting in less deterioration of thermally labile components. In addition, SFE with CO₂ can ensure that the active ingredients undergo minimal alterations, preserving their curative properties (Nasser et al., 1992; Oszagyan et al., 1996). Thus, SFE with CO₂ can be considered as an effective extraction method to obtain bio-actives.

Marine algae have intensively been studied for their vast diversity of physiological effects, including whitening, anti-inflammatory, anti-cancer, and anti-obesity activity (El Gamal, 2010; Li et al., 2011). *Ishige okamurae* is an edible brown alga found along the coast of Jeju Island in South Korea. Several

studies have reported the biological benefits of *I. okamurae* extracts, including its inhibitory effects on HIV (Ahn et al., 2006) and bacterial phospholipase A2 activity (Cho et al., 2008), anti-inflammatory effects (Kim et al., 2010, 2009), and anti-oxidant effects in free radical-mediated oxidative systems (Zou et al., 2008). However, the anti-inflammatory effects of the SFE from *I. okamurae* (SFEIO) remain unexplored. The main objectives of the current study were to evaluate the fatty acid composition of SFEIO and to investigate its anti-inflammatory activity in LPS-activated murine macrophage (RAW 264.7) cells.

MATERIAL AND METHODS

Reagents

Fetal bovine serum (FBS), Dulbecco's modified Eagle's medium (DMEM), antibiotics (penicillin and streptomycin were obtained from Invitrogen-Gibco (Grand Island, NY). Dimethyl sulfoxide (DMSO), lipopolysaccharide (LPS; from *Escherichia coli* strain) and phosphate buffered saline (PBS) were purchased from Sigma Chemical Co (St. Louis, MO, USA). The ELISA kits for TNF- α , PGE₂, IL-1 β , and IL-6 were obtained from BD Biosciences (San Diego, CA, USA) and R&D Systems, Inc. (St. Louis, MO, USA). Anti-bodies that includes: iNOS, COX-1, COX-2, I κ B- α , ERK, JNK, p38, p-ERK, p-JNK, p-p38 and β -actin were purchased from Cell Signaling Technology (Beverly, MA, USA). The secondary antibodies were obtained from Cell Signaling Technology.

Supercritical fluid extraction of *I. okamurae*

The marine alga *I. okamurae* was collected off the Jeju Island, South Korea in March 2011 and was identified by Dr. Dong-Sam Kim. The sample was dually washed using tap water to remove salt, epiphytes, and other debris attached to the surface, rinsed carefully in fresh water, and dried at room temperature for 2 weeks. The dried sample was ground into a powder prior to extraction. A supercritical extraction system

(ILSHIN Autoclave Co., Daejeon, Korea) with a 500 mL extraction cell was used to perform SFE on *I. okamurae*. The extraction cell was filled with 300 g of powdered *I. okamurae*. The extraction was conducted at 40° C for 2 h, providing an extraction pressure of 400 bars.

Fatty acid analysis

Fatty acid profile of the sample was determined using a modified fatty acid methyl ester (FAME) method (Morrison and Smith, 1964). Initially, 18.38 mg of the sample was weighed into a 30 mm × 110 mm Pyrex tubes fitted with a Teflon-lined cap. Undecanoic acid in dichloromethane was added as the internal standard. A volume of 1.5 mL of 0.5 N NaOH in methanol was introduced to the tubes, vortexed and incubated at 100° C for 5 min. After the incubation, 2 mL of boron trifluoride (14 %) in methanol was introduced and incubated at 100° C for 5 min. Tubes were then removed and cooled to room temperature before adding hexane (1 mL) to extract the FAMEs. A solution of saturated sodium chloride (5 mL) was added, and the mixture was shaken briefly and kept undisturbed allowing the phase separation. The hexane phase was then transferred to a conical tube. The tube remains were again partitioned with hexane (1 mL). The hexane phase was filtered through a silicon-treated Whatman filter paper to remove any residual water.

A Shimadzu GC-2010 plus gas chromatographic (GC) system, equipped with an SP-2560 fused silica capillary column (100 m × 0.25 mm i.d. and 0.2 µm film thickness) (Supelco, Inc., Bellefonte, PA, USA) coupled to a flame ionization detector (FID) was used to analyze the FAMEs. Helium was used as carrier gas at a continuous flow rate of 2.41 mL/min, given a head pressure of 403.1 kPa and a 1:100 split ratio. Both the injector and detector temperatures were held at 260° C. The oven temperature was increased from 100 to 240° C at a rate of 4° C/min and then held constant for 20 min. GC retention time and the fragmentation pat-

terns of FAMEs were used to identify the components. Quantitative measurements were taken using their response factors calculated by comparing the peak areas of the standards to the internal standard, undecanoic acid.

Cell culture

Murine macrophage RAW 264.7 cells were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA). The cells were maintained at 5×10^5 cells/mL in DMEM medium supplemented with 10 % FBS, Each 100 units/mL of antibiotics, penicillin and streptomycin were incubated at 37° C in a humidified atmosphere containing 5 % CO₂.

Determination of NO production

The nitrite concentration in the medium was measured as an indicator of NO production using the Griess reagent. For this, RAW 264.7 cells (2×10^5 cells/ml) were pre-incubated for 24 hours at 37° C following the treatment of SFEIO and LPS (1µg/ml), the quantity of nitrite accumulated in the culture medium was measured as an indicator of NO production. The evaluation was done by mixing equal volumes of culture medium and Griess reagent (1 % sulfanilamide in 5 % phosphoric acid and 0.1 % naphthylethylenediamine dihydrochloride in distilled water). The mixture was allowed to reach the equilibrium at room temperature for 10 min, and the absorbance was determined at 540 nm with a microplate reader. Fresh culture medium was used as the blank in every experiment. Nitrite levels in samples were determined using a standard sodium nitrite curve.

Lactic dehydrogenase (LDH) cytotoxicity assay

Sophocarpine-induced cytotoxicity was measured using the cytotoxicity detection kit (LDH) (Nanjing Jiancheng Co.). RAW 264.7 cells (1.5×10^5 cells/ml) were plated in 96 well plates and pre-incubated for 24 hours. Then the samples and LPS (1 µg/ml) were treated to the plates following an incubation

period of 24 h at 37° C. The medium was then collected, and the amount of LDH released by cells was determined using an assay kit according to the manufacturer's instructions. Finally, the absorbance of each well was measured at 490 nm using a microplate reader.

Determination of pro-inflammatory cytokines (TNF- α , IL-1 β , and IL-6) and prostaglandin E₂ (PGE₂) production

Macrophages were cultured in 24 well plates allowing 24 hours of post-incubation and treated with different concentrations of SFEIO (SFE from *I. okamurae*) followed by LPS (1 μ g/ml) and further incubated for another 24 hours at 37° C. Finally, the cytokines and PGE₂ concentrations in the culture medium were quantified using a competitive enzyme immunoassay kit (R&D Systems, Minneapolis, MN, USA) according to the manufacturer's instructions.

Immunoblotting

RAW 264.7 macrophages were cultured at a concentration of 5×10^5 cells per dish and incubated at 37° C for 24 h. Then the cells were treated with different concentrations of SFEIO (SFE from *I. okamurae*) and incubated for 24 h. The cells were lysed using a RIPA buffer [50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1 % Triton X-100, 1 % sodium deoxycholate, 0.1 % sodium dodecyl sulfate (SDS)]. Protein concentrations were determined using a Bio-Rad protein assay kit (Bio-Rad, CA, USA) using bovine serum albumin (BSA) as the calibration standard. Cell lysates were electrophoresed in SDS-polyacrylamide gels (8–12 %), and the separated proteins were transferred to PVDF membranes (Bio-Rad). The immunoblot was kept in the blocking solution (Tris-buffered saline/Tween 20, TBST) containing 5 % skim milk (w/v) for 3 h under gentle shaking conditions at room temperature. Then the membranes were incubated with primary antibodies, iNOS, COX-2, I κ B- α , p-p50, p-ERK, ERK, p-JNK and JNK (cell signaling; 1:1000) in 5 % BSA in

TBST for 24h at 4° C. After the removal of the primary antibodies, the membranes were washed three times with TBST buffer at room temperature, and incubated with peroxidase-conjugated secondary antibodies in 5 % BSA in TBST (diluted 1:5000; Cell signaling) for 2 h at room temperature. Following the addition, signals were developed using ECL western blotting detection kit exposed to Biorad Chemidac system.

Statistical analysis

All data in this study are expressed as means \pm S.D. Significant differences among the groups were determined using the unpaired Student's *t*-test. A value of **p* < 0.05 was considered to be statistically significant.

RESULTS

Fatty acid profile of SFEIO

Table 1, summarize the results of the fatty acid analysis for the SFEIO and the Figure 1 represents the corresponding chromatograms for the fatty acid composition in SFEIO and the used standard. A total of 37 fatty acids were investigated, and 17 of them were identified in SFEIO. Palmitic acid (30.2 %) was the most abundant constituent in SFEIO, followed by linoleic acid (23.1 %) and oleic acid (16.9 %). Total fatty acid content in SFEIO calculated as the sum of the individual fatty acids, was 728.4 mg/g extract wt; the total saturated and the unsaturated fatty acid content was 292.4 and 436.0 mg/g extract wt, respectively contributing to 40.1 % and 59.9 % of the total fatty acid content.

Effect of SFEIO on production of NO and PGE₂ in LPS-stimulated RAW 264.7 cells

The effect of SFEIO on LPS-stimulated NO and PGE₂ production in RAW 264.7 cells were assessed by treating the cells with LPS except for the control following the treatment of SFEIO (25, 50, or 100 μ g/mL). Nitrite and PGE₂ levels in unstimulated RAW 264.7 cells were negligible; however,

Table 1: Fatty acid composition in SFEIO

No	Fatty acid name		Abundance (%)	mg/g extract
1	C4:0	Butyric acid	-	-
2	C6:0	Caproic acid	-	-
3	C8:0	Caprylic acid	-	-
4	C10:0	Capric acid	-	-
5	C11:0	Undecylic acid	-	-
6	C12:0	Lauric acid	-	-
7	C13:0	Tridecylic acid	-	-
8	C14:0	Myristic acid	7.0	51.2
9	C14:1	Myristoleic acid	-	-
10	C15:0	Pentadecylic acid	0.6	4.6
11	C15:1	<i>Cis</i> -10-pentadecenoic acid	-	-
12	C16:0	Palmitic acid	30.2	220.2
13	C16:1	Palmitoleic acid	2.4	17.4
14	C17:0	Heptadecanoic acid	0.3	2.2
15	C17:1	<i>Cis</i> -10-heptadecenoic acid	-	-
16	C18:0	Stearic acid	2.0	14.2
17	C18:1(n-9), <i>trans</i>	Elaidic acid	-	-
18	C18:1(n-9), <i>cis</i>	Oleic acid	16.9	123.0
19	C18:2(n-6), <i>trans</i>	Linolelaidic acid	-	-
20	C18:2(n-6), <i>cis</i>	Linoleic acid	23.1	168.0
21	C20:0	Arachidic acid	0.3	2.3
22	C18:3(n-6)	Gamma-linolenic acid	0.6	4.2
23	C20:1(n-9)	Eicosenoic acid	-	-
24	C18:3(n-3)	Alpha-linolenic acid	2.4	17.6
25	C21:0	Heneicosylic acid	-	-
26	C20:2(n-6)	Eicosadienoic acid	1.7	12.4
27	C22:0	Behenic acid	-	-
28	C20:3(n-6)	Dihomo-gamma-linolenic acid	0.4	3.0
29	C22:1(n-9)	Erucic acid	-	-
30	C20:3(n-3)	Eicosatrienoic acid	0.3	2.1
31	C23:0	Tricosylic acid	-	-
32	C20:4(n-6)	Arachidonic acid	7.9	57.7
33	C22:2(n-6)	Docosadienoic acid	0.3	2.4
34	C24:0	Lignoceric acid	-	-
35	C20:5(n-3)	Eicosapentaenoic acid	3.6	25.9
36	C24:1(n-9)	Nervonic acid	-	-
37	C22:6(n-3)	Docosahexaenoic acid	-	-
Total saturated fatty acid			40.1	292.4
Total unsaturated fatty acid			59.9	436.0
Total fatty acid^a			100	728.4

^aAbundance % is the proportion of individual fatty acid to the sum of the total fatty acids identified and determined through GC-MS.




Figure 1: Chromatogram of the FAME of the **(A)** standard (C4:0-C22:6 methyl esters) and **(B)** SFEIO

their levels were increased substantially following the LPS stimulation (Figure 2). Treatment of RAW 264.7 cells with SFEIO significantly and dose-dependently inhibited LPS-stimulated NO and PGE₂ production (Figure 2). To examine whether the inhibitory effects of SFEIO on production of pro-inflammatory mediators in LPS-stimulated RAW 264.7 cells were due to cytotoxicity (Figure 2A). Based on the results, SFEIO did not induce cytotoxicity in RAW 264.7 cells at the evaluated concentrations.

Effects of SFEIO on iNOS and COX-2 protein expression in LPS-stimulated RAW 264.7 cells

iNOS and COX-2 are two of the most important signaling molecules that regulate inflammatory reactions. Western blot analysis was used to determine the effects of SFEIO on iNOS and COX-2 protein expression in RAW 264.7 cells upon treatment with different concentrations of SFEIO (25, 50, or 100 µg/mL) as SFEIO was found to inhibit NO and PGE₂ production in LPS-stimulated RAW 264.7 cells. According to the results, iNOS and COX-2 protein expression were almost undetectable in unstimulated

RAW 264.7 cells. Whereas, treatment with 1.0 µg/mL LPS significantly increased the iNOS and COX-2 protein expression. However, the treatment of SFEIO significantly and dose-dependently downregulated the iNOS and COX-2 protein expression in LPS-stimulated RAW 264.7 cells (Figure 3). Therefore, the inhibition of NO and PGE₂ production in LPS-stimulated RAW 264.7 cells is attributed to the inhibition of iNOS and COX-2 expression.

Effects of SFEIO on pro-inflammatory cytokine expression in LPS-stimulated RAW 264.7 cells

Tumor necrosis factor (TNF)-α, interleukin (IL)-1β, and IL-6 are some of the pro-inflammatory cytokines that play important roles in immune responses variety of inflammatory stimuli (Bertolini et al., 2001; Lind, 2003). Therefore, the inhibitory effects of SFEIO on LPS-stimulated TNF-α, IL-1β, and IL-6 production were dually examined to identify the expression levels. These levels were measured in culture supernatants using enzyme-linked immunosorbent assay (ELISA) kits. Treatment of RAW 264.7 cells with LPS significantly up-regulated the cytokine production, compared to the non-treated group. However, TNF-α, IL-1β, and IL-6 levels were significantly and dose-dependently reduced relative to the positive control treated with LPS (Figure 4).

Effect of SFEIO on LPS-induced NF-κB and mitogen-activated protein kinase (MAPK) activation

NF-κB plays an important role in the transcription of pro-inflammatory mediators following LPS stimulation (Henkel et al., 1993; Rhee et al., 2007). Therefore, western blot analysis was performed to assess the effect of SFEIO on IκBα and p50 NF-κB phosphorylation. As shown in Figure 4A, treatment with LPS induced IκBα degradation, which was significantly blocked by SFEIO. SFEIO also dose-dependently inhibited LPS-induced p50 NF-κB activation (Figure 5A).




Figure 2: Effects of SFEIO on (A) NO and (B) PGE₂ production in LPS-induced RAW 264.7 cells. Cells stimulated with LPS (1 µg/ml) in the presence of SFEIO (25, 50, and 100 µg/ml) for 24 h at 37° C. (A) Cytotoxicity was determined using the LDH method. Values are expressed as means ± S.D. of triplicate experiments. **P* < 0.05 indicate significant differences from the LPS-stimulated group.




Figure 3: Effects of SFEIO on LPS-induced iNOS and COX-2 protein expressions in RAW 264.7 cells. Cells stimulated with LPS (1 µg/ml) in the presence of SFEIO (25, 50, and 100 µg/ml) for 24 h at 37° C. iNOS and COX-2 protein level were determined via Western blotting.




Figure 4: Inhibitory effects of SFEIO on the pro-inflammatory cytokine production in RAW 264.7 cells. The production of (A) TNF- α , (B) IL-6, and (C) IL-1 β were assayed in the culture medium of cells stimulated with LPS (1 μ g/ml) for 24 h in the presence of SFEIO (25, 50, and 100 μ g/ml). Supernatants were collected, and the TNF- α , IL-6, and IL-1 β concentration in the supernatants were determined by ELISA. Values are expressed as means \pm S.D. of triplicate experiments. * $P < 0.05$.

MAPK are known to regulate the expression of inflammatory enzymes and cytokines (Hommes et al., 2003). To determine whether the inhibition of inflammation was mediated via the MAPK pathway, phosphorylation of p38, extracellular-signaling related kinase 1/2 (ERK1/2), and c-Jun N-terminal kinase (JNK) in LPS-induced cells were examined by western blot. As shown in Figure 4B, LPS-stimulation markedly increased p38, ERK1/2, and JNK phosphorylation. However, SFEIO significantly inhibited LPS-stimulated ERK and JNK phosphorylation. Importantly, the levels of non-phosphorylated MAPK isoforms did not vary significantly among groups (Figure 5B). In contrast, SFEIO treatment did not affect the LPS-stimulated p38 protein phosphorylation (data not shown).




Figure 5: Inhibitory effects of SFEIO on LPS-induced phosphorylation of (A) I κ B- α , NF- κ B p50, (B) JNK, and ERK. Cells were treated for 15 min with LPS (1 μ g/ml) alone or with LPS (1 μ g/ml) coupled with different concentrations (025, 50, and 100 μ g/ml) of SFEIO. Cell lysates were extracted, and protein levels of p- I κ B- α , p-p50 p-ERK, ERK, p-JNK, and JNK were analyzed by Western blot.

DISCUSSION

SFE has several advantages over conventional solvent extraction, as it shows increased selectivity, decreased time consumption, and it is environmentally friendly (Shu et al., 2004). However, *I. okamurae* has not previously been extracted using SFE for bioactivity screenings. Thus, we investigated the anti-inflammatory effects and fatty acid constituents of *I. okamurae* extracts by using SFE technology.

Fatty acids from seaweeds can be obtained by using SFE as extensively reported (Cheung, 1999; Herrero et al., 2006). Therefore, we first measured the fatty acids composition of SFEIO. The major fatty acid components in the extract were palmitic acid (30.2 %), linoleic acid (23.1 %) and oleic acid (16.9 %). Unsaturated fatty acids contributed to 59.9 % of the total fatty acid content. Fatty acids such as phospholipids play a key role in maintaining the structural integrity of the cell membrane and play important roles in maintaining various cellular functions including metabolism and immune responses (Basu et al., 2006; Cabral, 2005; Natali et al., 2007). The dietary intake of mono- and polyunsaturated fatty acids is associated with reduced risk of cardiovascular diseases including hypertension and atherosclerosis (Basu et al., 2006). Polyunsaturated linoleic acid inhibits COX-1 and COX-2 in vitro (Ciaraldi et al., 2002; Ringbom et al., 2001) and decreases the expression levels of NF- κ B and TNF- α genes in human THP-1 monocytic cells, leading to a decreased production of IL-6, IL-1 β and TNF- α levels (Zhao et al., 2005). Recently, the monounsaturated oleic acid was found to inhibit LPS-induced iNOS and COX-2 expression through blocking p38 and NF- κ B signaling pathways in BV2 murine microglia (Oh et al., 2009). Palmitic acid has been shown to trigger inflammation in cultured human macrophages (Laine et al., 2007) and increasing TNF- α and IL-6 production via p38 and ERK activation in astrocytes cells (Gupta et al., 2012). However, ω -3 fatty acids could act in a dose-dependent manner preventing the palmitic

acid-induced inflammatory responses in astrocytes (Gupta et al., 2012). Moreover, oleic acid is able to reverse the saturated fatty acid palmitate-induced insulin resistance and inflammation in skeletal muscle cells (Coll et al., 2008). Accordingly, we expected the SFEIO, which contained unsaturated fatty acids, to exhibit anti-inflammatory activities.

Macrophages play a key role in immune responses during inflammation. Activation of macrophages by LPS releases inflammatory mediators, including NO and PGE₂, which are generated via the oxidation of the terminal guanidine nitrogen of L-arginine by iNOS and the conversion of arachidonic acid by COX-2 (Posadas et al., 2000). Both NO and PGE₂ are involved in the progression of inflammation and carcinogenesis (Claria, 2003). IL-1 β , IL-6, and TNF- α are some of the major pro-inflammatory cytokines, which are produced by monocytes and macrophages which play a crucial role in the progression of the initial steps in the inflammatory response (Tao et al., 2009; Young et al., 2002). Thus, reducing the levels of NO, PGE₂, IL-1 β , IL-6, and TNF- α could be an effective strategy to inhibit inflammation. Current results suggest that SFEIO could inhibit the production of NO and PGE₂ by the suppression of iNOS and COX-2 expression. Additionally, SFEIO reduced the production of IL-1 β , IL-6, and TNF- α in LPS-induced RAW 264.7 cells.

NF- κ B-mediated regulation of pro-inflammatory gene expression is controlled by transferases belonging to the I κ B family. The transcription factor NF- κ B could regulate the expression of pro-inflammatory mediators, including COX-2, iNOS, IL-1 β , IL-6, and TNF- α (Kim et al., 2010; Rhee et al., 2007). indicate that the degradation and phosphorylation of I κ B, a cytoplasmic inhibitor of NF- κ B, could induce inflammatory stimuli, resulting in the translocation of the activated p65/p50 heterodimer from the cytoplasm to the nucleus. Current results indicate that SFEIO suppression the LPS induced I κ B- α phosphorylation and degradation, and the activation of NF- κ B p50. These

results reveal that the effects of SFEIO on inflammatory mediators and cytokine production are mediated via the inhibition of NF- κ B pathway.

Additionally, SFEIO could attenuate LPS-stimulated phosphorylation of JNKs and ERK, whereas it did not affect p38 MAPK. MAPKs include ERKs, JNKs, and p38 kinase (Shao et al., 2013), and the phosphorylation of these proteins promotes the pro-inflammatory cytokine production in LPS-induced macrophages (Kim et al., 2013; Yoon et al., 2012). In particular, activation of ERK and JNK could regulate the responses in LPS-stimulated macrophages, inducing increased pro-inflammatory cytokine and iNOS production (Ajizian et al., 1999; Uto et al., 2005). These results demonstrate that the anti-inflammatory effect of SFEIO could down-regulate LPS-stimulated phosphorylation of JNK and ERK in RAW 264.7 cells.

In conclusion, the present study revealed that SFEIO treatment inhibited LPS-stimulated NO and PGE₂ production in macrophages by suppressing iNOS and COX-2 expression, respectively. SFEIO also significantly inhibited the release of IL-1 β , IL-6, and TNF- α in LPS-stimulated RAW 264.7 cells. These effects occurred due to inhibition of NF- κ B signaling and JNK and ERK phosphorylation. These findings indicate that SFEIO may have potential anti-inflammatory activity.

Acknowledgment

This research was supported by the project (C36290) offered by the Korea Basic Science Institute. This study was supported by a research grant from the Marine Biotechnology Program funded by the Ministry of Oceans and Fisheries of Korean Government (PM59122), and was partially supported by a research grant funded by the Korea Institute of Ocean Science & Technology (PE99411).

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Ahn MJ, Yoon KD, Kim CY, Kim JH, Shin CG, Kim J. Inhibitory activity on HIV-1 reverse transcriptase and integrase of a carmalol derivative from a brown Alga, *Ishige okamurae*. *Phytother Res.* 2006;20:711-3.
- Ajizian SJ, English BK, Meals EA. Specific inhibitors of p38 and extracellular signal-regulated kinase mitogen-activated protein kinase pathways block inducible nitric oxide synthase and tumor necrosis factor accumulation in murine macrophages stimulated with lipopolysaccharide and interferon-gamma. *J Infect Dis.* 1999;179:939-44.
- Basu A, Devaraj S, Jialal I. Dietary factors that promote or retard inflammation. *Arterioscler Thromb Vasc Biol.* 2006;26:995-1001.
- Bertolini A, Ottani A, Sandrini M. Dual acting anti-inflammatory drugs: a reappraisal. *Pharmacol Res.* 2001;44:437-50.
- Cabral GA. Lipids as bioeffectors in the immune system. *Life Sci.* 2005;77:1699-710.
- Cheung PCK. Temperature and pressure effects on supercritical carbon dioxide extraction of n-3 fatty acids from red seaweed. *Food Chem.* 1999;65:399-403.
- Cho JY, Gyawali YP, Ahn SH, Khan MN, Kong IS, Hong, YK. A methoxylated fatty acid isolated from the brown seaweed *Ishige okamurae* inhibits bacterial phospholipase A2. *Phytother Res.* 2008;22:1070-4.
- Ciaraldi TP, Cha BS, Park KS, Carter L, Mudaliar SR, Henry RR. Free fatty acid metabolism in human skeletal muscle is regulated by PPARgamma and RXR agonists. *Ann N Y Acad Sci.* 2002;967:66-70.
- Claria J. Cyclooxygenase-2 biology. *Curr Pharm Des.* 2003;9:2177-90.
- Coll T, Eyre E, Rodriguez-Calvo R, Palomer X, Sanchez RM, Merlos M, et al. Oleate reverses palmitate-induced insulin resistance and inflammation in skeletal muscle cells. *J Biol Chem.* 2008;283:11107-16.
- El Gamal AA. Biological importance of marine algae. *Saudi Pharm J.* 2010;18:1-25.
- Gupta S, Knight AG, Gupta S, Keller JN, Bruce-Keller AJ. Saturated long-chain fatty acids activate inflammatory signaling in astrocytes. *J Neurochem.* 2012;120:1060-71.

- Henkel T, Machleidt T, Alkalay I, Kronke M, Ben-Neriah Y, Baeuerle PA. Rapid proteolysis of I kappa B-alpha is necessary for activation of transcription factor NF-kappa B. *Nature*. 1993;365:182-5.
- Herrero M, Cifuentes A, Ibañez E. Sub- and supercritical fluid extraction of functional ingredients from different natural sources: Plants, food-by-products, algae and microalgae: A review. *Food Chem*. 2006; 98:136-48.
- Hommel DW, Peppelenbosch MP, van Deventer SJ. Mitogen activated protein (MAP) kinase signal transduction pathways and novel anti-inflammatory targets. *Gut*. 2003;52:144-51.
- Kim KN, Heo SJ, Yoon WJ, Kang SM, Ahn G, Yi TH, et al. Fucoxanthin inhibits the inflammatory response by suppressing the activation of NF-kappaB and MAPKs in lipopolysaccharide-induced RAW 264.7 macrophages. *Eur J Pharmacol*. 2010;649:369-75.
- Kim KN, Ko YJ, Yang HM, Ham YM, Roh SW, Jeon YJ, et al. Anti-inflammatory effect of essential oil and its constituents from fingered citron (*Citrus medica* L. var. *sarcodactylis*) through blocking JNK, ERK and NF-kappaB signaling pathways in LPS-activated RAW 264.7 cells. *Food Chem Toxicol*. 2013;57:126-31.
- Kim MM, Rajapakse N, Kim SK. Anti-inflammatory effect of *Ishige okamurae* ethanolic extract via inhibition of NF- κ B transcription factor in RAW 264.7 cells. *Phytother Res*. 2009;23:628-34.
- Korhonen R, Lahti A, Kankaanranta H, Moilanen E. Nitric oxide production and signaling in inflammation. *Curr Drug Targets Inflamm Allergy*. 2005;4: 471-9.
- Laine PS, Schwartz EA, Wang Y, Zhang WY, Karnik SK, Musi N, et al. Palmitic acid induces IP-10 expression in human macrophages via NF-kappaB activation. *Biochem Biophys Res Commun*. 2007; 358:150-5.
- Li Y-X, Wijesekara I, Li Y, Kim S-K. Phlorotannins as bioactive agents from brown algae. *Process Biochem*. 2011;46:2219-24.
- Lind L. Circulating markers of inflammation and atherosclerosis. *Atherosclerosis*. 2003;169:203-14.
- Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008;454:428-35.
- Morrison WR, Smith LM. Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride-methanol. *J Lipid Res*. 1964;5:600-8.
- Nasser W, Reverchon S, Robert-Baudouy J. Purification and functional characterization of the KdGR protein, a major repressor of pectinolysis genes of *Erwinia chrysanthemi*. *Mol Microbiol*. 1992;6:257-65.
- Natali F, Siculella L, Salvati S, Gnoni GV. Oleic acid is a potent inhibitor of fatty acid and cholesterol synthesis in C6 glioma cells. *J Lipid Res*. 2007;48: 1966-75.
- Oh YT, Lee JY, Lee J, Kim H, Yoon KS, Choe W, et al. Oleic acid reduces lipopolysaccharide-induced expression of iNOS and COX-2 in BV2 murine microglial cells: possible involvement of reactive oxygen species, p38 MAPK, and IKK/NF-kappaB signaling pathways. *Neurosci Lett*. 2009;464:93-7.
- Oszagyan M, Simándi B, Sawinsky J, Kéry Á, Lemberkovic E, Fekete J. Supercritical fluid extraction of volatile compounds from Lavandin and Thyme. *Flavour Frag J*. 1996;11:157-65.
- Posadas I, Terencio MC, Guillen I, Ferrandiz ML, Coloma J, Paya M, et al. Co-regulation between cyclo-oxygenase-2 and inducible nitric oxide synthase expression in the time-course of murine inflammation. *Naunyn Schmiedeberg's Arch Pharmacol*. 2000;361: 98-106.
- Rhee JW, Lee KW, Kim D, Lee Y, Jeon OH, Kwon HJ, et al. NF-kappaB-dependent regulation of matrix metalloproteinase-9 gene expression by lipopolysaccharide in a macrophage cell line RAW 264.7. *J Biochem Mol Biol*. 2007;40:88-94.
- Ringbom T, Huss U, Stenholm Å, Flock S, Skattebøl L, Perera P, et al. COX-2 Inhibitory effects of naturally occurring and modified fatty acids. *J Nat Prod*. 2001;64:745-9.
- Shao J, Li Y, Wang Z, Xiao M, Yin P, Lu Y, et al. 7b, a novel naphthalimide derivative, exhibited anti-inflammatory effects via targeted-inhibiting TAK1 following down-regulation of ERK1/2- and p38 MAPK-mediated activation of NF-kappaB in LPS-stimulated RAW 264.7 macrophages. *Int Immunopharmacol*. 2013;17:216-28.
- Shu XS, Gao ZH, Yang XL. Supercritical fluid extraction of sapogenins from tubers of *Smilax china*. *Fitoterapia*. 2004;75:656-61.
- Tao JY, Zheng GH, Zhao L, Wu JG, Zhang XY, Zhang SL, et al. Anti-inflammatory effects of ethyl acetate fraction from *Melilotus suaveolens* Ledeb on LPS-stimulated RAW 264.7 cells. *J Ethnopharmacol*. 2009;123:97-105.

Uto T, Fujii M, Hou DX. 6-(Methylsulfinyl)hexyl isothiocyanate suppresses inducible nitric oxide synthase expression through the inhibition of Janus kinase 2-mediated JNK pathway in lipopolysaccharide-activated murine macrophages. *Biochem Pharmacol.* 2005;70:1211-21.

Yoon WJ, Heo SJ, Han SC, Lee HJ, Kang GJ, Kang HK, et al. Anti-inflammatory effect of sargachromanol G isolated from *Sargassum siliquastrum* in RAW 264.7 cells. *Arch Pharm Res.* 2012;35:1421-30.

Young JL, Libby P, Schonbeck U. Cytokines in the pathogenesis of atherosclerosis. *Thromb Haemost.* 2002;88:554-67.

Zhao G, Etherton TD, Martin KR, Vanden Heuvel JP, Gillies PJ, West SG, et al. Anti-inflammatory effects of polyunsaturated fatty acids in THP-1 cells. *Biochem Biophys Res Commun.* 2005;336:909-17.

Zou Y, Qian ZJ, Li Y, Kim MM, Lee SH, Kim SK. Antioxidant effects of phlorotannins isolated from *Ishige okamurae* in free radical mediated oxidative systems. *J Agric Food Chem.* 2008;56:7001-9.